

LIST OF FIGURES

Figure No.	Caption	Page No.
1.1	Structure of zeolite Y	03
1.2	(A) Basic building blocks of zeolite framework, (B) Linkage between sodalite cage and supercage of zeolite-Y	04
1.3	Different approaches for the synthesis of metal complex inside the cavity of zeolite Y. (A) Flexible ligand (B) Template synthesis method and (C) Zeolite synthesis method.	08
1.4	(A) Cytochrome P450 bioenzymes. (B) Co(II) phthalocyanine complex inside zeolite-Y	09
1.5	Magnetic moments of (a) $[\text{Co}(\text{bpy})_3](\text{ClO}_4)_2$ and (b) $[\text{Co}(\text{bpy})_3]^{2+}$ encapsulated in zeolitY	15
1.6	Molecular graphics representation of a square-planar Co phthalocyanine molecule encapsulated within the supercage of zeolite-Y	16
1.7	Total potential energy for CoPc encapsulated in zeolite-Y as a function of the average rms displacement of the atoms of the Pc ring from planar geometry (C_{av}). The saddle-type distortion is shown in the inset. (B) Temperature variation of magnetic susceptibilities of (a) CoPc in free state (b) encapsulated in MCM-41 and in (c) zeolite Y	17
1.8	Schematic views of (A and B) $[\text{Fe}(\text{phen})_3]^{2+}$ in the supercage and (C) one of the three ligands facing the 12-membered ring of the cage	19
1.9	Different perspective of DFT optimized structure of (left) the free and (right) the encapsulated nickel Schiff-base complex obtained at the B3LYP level	20
1.10	Cyclic voltammogram of (A) CoPhen (neat) and (B) CoPhenY	20
1.11	(A-B) UV-Vis spectra of (a) <i>N,N'</i> -bis(salicylidene)ethylenediamine (Salen), (b) <i>N,N'</i> -disalicylidene-1,2-phenylenediamine (Salophen), (c) neat Fe-Salen complex, and (d) neat Fe-Salophen complex And (C) DR spectra of ion exchanged zeolites (a) Fe-LiY, (b) Fe-NaY, (c) Fe-KY	21
1.12	2,2-bipyridine and 1,10-phenanthroline with iron(III) and nickel(II) in a zeolite Y	22

LIST OF FIGURES

1.13	Schematic representation with the molecular dimension (end-to-end distance) of M (salen) complexes. (where M = Ni, Pd, Co, Cu)	23
2.1	Photograph of powder X-ray diffractometer (RIGAKU MiniFlex II)	36
2.2	Photograph of Scanning Electron Microscope Zeiss EVO 40	37
2.3	2.3: Photograph of X-ray photoelectron spectrophotometer (Omicron EA 125)	38
2.4	Photograph of the Fourier Transform Infrared (FT-IR) spectrometer..	38
2.5	Photograph of the UV-Vis spectrophotometers (a) Shimadzu with model UV-2450 (b) JASCO with model V-650	40
2.6	Photograph of Atomic Absorption Spectrophotometer (Shimadzu AA-7000).	40
2.7	Photograph of SQUID magnetometer Quantum Design MPMS XL Ever Cool.	41
2.8	Photograph of GC-Shimadzu-2014.	42
3A.1	(A) Powder XRD pattern of (a) pure zeolite-Y, (b) Ni exchanged zeolite-Y, (c) Encapsulated NiL3 in zeolite Y, (B) The XRD patterns of (a) Al-MCM-41, (b) Ni-Al-MCM-41 and (c) Encapsulated NiL3 in MCM-41 at the range of ($0 < 2\theta < 4$).	47
3A.2	SE micrograph of encapsulated NiL3 in zeolite Y, (A) before Soxhlet extraction, (B) after Soxhlet extraction.	49
3A.3	(A) FTIR spectra in the range of 500 cm^{-1} to 4000 cm^{-1} of (a) pure zeolite Y, (b) encapsulated NiL3 in zeolite Y (B) The FTIR spectra in the range of 600 cm^{-1} to 2000 cm^{-1} (a) pure zeolite Y, (b) encapsulated NiL3 in zeolite Y, (c) “neat” NiL3 complex. (C) FTIR spectra in the range of 500 cm^{-1} to 4000 cm^{-1} of (a) pure MCM-41, (b) encapsulated NiL3 in MCM-41	50
3A.4	(a) NiL3 molecule (singlet) free state (b) NiL3 molecule (singlet) is encapsulated and extracted (c) NiL3 molecule (triplet) is encapsulated and extracted. The atoms marked by the square define the C-Ni-C bond angle. (d) The zeolite framework used for theoretical studies (e) NiL3molecule (triplet)	52

LIST OF FIGURES

	is encapsulated within the zeolite pore. Closest non-bonding distance between NiL3 and the zeolite framework is $\sim 3.1 \text{ \AA}$.	
3.5	The structure and important molecular orbitals for NiL3. The frontier orbitals with clear metal orbital characters are shown	54
3A.6	(a) The molecular orbitals of NiL3 in free state, including the HOMO, LUMO. The orbitals lower in energy than HOMO are marked as H-n ($n = 0$ being HOMO) and LUMO+n ($n = 0$ being LUMO). (b) The molecular orbitals of NiL3 in encapsulated singlet state, including the HOMO, LUMO. Corresponding molecular structures are shown for reference.	55
3A.7	The molecular orbitals of NiL3 in encapsulated triplet state, including the HOMO, LUMO. The orbitals lower in energy than HOMO are marked as HOMO-n ($n = 0$ being HOMO) and LUMO+n ($n = 0$ being LUMO). Corresponding molecular structure is shown for reference.	57
3A.8	Experimental and DFT simulated UV-Vis spectra of (A) DFT simulated singlet state (black vertical plot) and experimental encapsulated NiL3 in zeolite Y, (red line plot) (B) DFT simulated triplet state (black vertical plot) and experimental encapsulated NiL3 in zeolite Y, (red line plot).	57
3A.9	UV-Vis spectra of (a) L3 in chloroform (b) TD- DFT simulated UV-Vis spectra of L3.	60
3A.10	(A) The solid state UV-Vis spectra of (a) NiL3 (b) encapsulated NiL3 in zeolite, (c) NiL3 in chloroform (B) UV-Visible spectra of (a) "neat" NiL3 in chloroform (b) encapsulated NiL3 in complex in MCM-41.	61
3A.11	The $1/x-x_0$ Vs T graph of (a) NiL3 complex in zeolite Y and (b) NiL3 complex in MCM-41.	63
3B.1	Schematic structure for the Ni-salen complexes in free state.	70
3B.2	XRD patterns of (a) pure zeolite Y, (b) Ni-exchanged Y, (c) NiL1-Y, (d) NiL4-Y and (e) NiL6-Y complexes.	72
3B.3	SEM images of pure (a) zeolite Y, and (b) NiL1-Y, (c) NiL6-Y after Soxhlet extraction.	72

LIST OF FIGURES

3B.4	(A) XPS survey spectra (a) NiL1, (b) NiL1-Y, and (c) NiL6-Y, and high resolution XPS spectra of Ni (2p) for (B) NiL1 and (C) NiL1-Y. (Black colored graphs are experimental data and green and blue colored peaks are peak fitted data)	74
3B.5	High resolution XPS spectra of (A) C (1s), (B) N (1s) and (C) O (1s) for NiL1. (Black colored graphs are experimental data and green and blue colored peaks are peak fitted data)	74
3B.6	High resolution XPS spectra of (A) C (1s), (B) N (1s) and (C) O (1s) for NiL1-Y. (Black colored graphs are experimental data and green and blue colored peaks are peak fitted data)	74
3B.7	(A) FTIR spectra of Ligands (a) L1, (b) L4 and (c) L6. (B) (A) FTIR spectra of free state complexes (a) NiL1, (b) NiL4 and (c) NiL6. (C) FTIR spectra of (a) pure zeolite Y, (b) NiL1-Y, (c) NiL4-Y and (d) NiL6-Y. (D) Enlarged view FTIR spectra of (a) pure zeolite Y, (b) NiL1-Y, (c) NiL4-Y and (d) NiL6-Y.	76
3B.8	(A). The solid state UV-Vis spectra of “neat” complex (a) NiL1, (b) NiL4 and (c) NiL6. (B) The solid state UV-Vis spectra of encapsulated complexes in zeolite Y, (a) Pure zeolite Y, (b) NiL1-Y, (c) NiL4-Y and (d) NiL6-Y.	78
3B.9	Molar susceptibility vs. temperature plots in the range of 5K-300K of the complexes in free and encapsulated states (a) NiL1, (b) NiL6, (c) NiL4, (d) NiL1-Y, (e) NiL4-Y and (f) NiL6-Y.	80
3B.10	Schematic structure for the Ni-salen complex in free state. b. Optimized structure of NiL6 (C atoms in green) encapsulated within zeolite supercage, two views are shown. Optimized structures of (c) NiL1, (d) NiL4 and (e) NiL6 in neat (singlet) state and encapsulated and extracted (triplet) states are shown.	82
3B.11	Comparative UV-Visible data of (A-B) NiL1, (C-D) NiL4 and (E-F) NiL6 in free and encapsulated state, experimental data are represented by red line and TD-simulated data are represented by blue vertical line.	85
3C.1	Comparative molecular dimensions of NiL1 and NiL2 complexes.	93
3C.2	Powder XRD patterns of (a) pure zeolite-Y (b) Ni exchanged zeolite-Y (c) NiL2-Y	94

LIST OF FIGURES

3C.3	SEM images of (a) pure zeolite Y, (b) NiL2-Y after Soxhlet extraction.	96
3C.4	XPS spectra of NiL2-Y complex (A) survey spectra, High resolution XPS spectra of (B) nickel (2p), (C) carbon (1s), (D) nitrogen (1s) and (E) oxygen (1s). (Black colored graphs are experimental data and green and blue colored peaks are peak fitted data)	97
3C.5	(A) FTIR spectra in the range of 500 cm ⁻¹ to 4000 cm ⁻¹ of (a) NiL2 (b) NiL2-Y, and (c) pure zeolite Y. (B) magnified view of IR data of (a) NiL2 (b) NiL2-Y, and (c) pure zeolite Y.	98
3C.6	(A) UV-Vis spectra in chloroform in the range of 190-700 nm of (a) L2 and (b) NiL2. (B) Solid state UV-Vis spectra in the range of 190-700 nm of (a) NiL1 (b) NiL2 (c) NiL2-Y and (d) NiL1-Y.	100
3C.7	(a) NiL1 (singlet) free state (b) NiL2 free singlet state. The atoms marked by the squares define the C-Ni-C bond angle in different molecules. (c) The zeolite framework used for theoretical studies. (d) The deformation of encapsulated Ni-salen (bottom) shown with respect to its neat form (top). (e) The deformation of NiL2-Y (bottom) shown with respect to its neat form (top). (f) NiL2-Y molecule (triplet)	102
3C.8	Comparative UV-Visible data of (A) NiL1 (B) NiL1-Y (C) NiL2 and (D) NiL2-Y experimental data are represented by red line and TD-simulated data are represented by black vertical line.	105
3C.10	Molar susceptibility vs temperature plots in the range of 4K-300K of (a) NiL1 (b) NiL1-Y and (c) NiL2-Y.	106
4.1	SEM images (a) zeolite Y, (b) Pd-Y, (c) PdL3-Y (before Soxhlet extraction) (d) PdL3-Y (after Soxhlet extraction), (e) PdL4-Y, (f) PdL6-Y.	114
4.2	XRD pattern of (a) Pure zeolite-Y, (b) Pd-exchanged zeolite -Y, (c) PdL1-Y, (d) PdL3-Y, (e) PdL4-Y, (f) PdL5-Y, and (g) PdL6-Y.	115
4.3	High resolution XPS signals of (A) Pd (3d), (B) C (1s), (C) N (1s) and (D) O (1s) for PdL1 Complex. (Black colored graphs are experimental data and green and blue colored peaks are peak fitted data)	116
4.4	High resolution XPS signals of (A) Pd (3d), (B) C (1s), (C) N (1s) and (D) O (1s) for PdL1-Y Complex. (Black colored graphs are experimental data and green and blue colored peaks are peak fitted data)	116
4.5	High resolution XPS signals of (a) Pd (3d), (b) C (1s), (c) N (1s) and (d) O (1s) for PdL3-Y Complex. (Black colored graphs are experimental data and green and blue colored peaks are peak fitted data)	117
4.6	High resolution XPS signals of (A) Pd (3d), (B) C (1s), (C) N (1s) and (D) O	118

LIST OF FIGURES

	(1s) for PdL6-Y Complex. (Black colored graphs are experimental data and green and blue colored peaks are peak fitted data)	
4.7	(A) FTIR spectra of free state palladium salen complexes (a) PdL1, (b) PdL3, (c) PdL4, (d) PdL5, and (e) PdL6. (B) FTIR spectra of encapsulated palladium salen complexes in zeolite Y (a) Pure zeolite Y, (b) PdL1-Y, (c) PdL3-Y, (d) PdL4-Y, (e) PdL5-Y and (f) PdL6-Y.	119
4.8	(A) UV-Vis spectra of free state complexes in chloroform, (a) PdL3, (b) PdL5, (c) PdL6 (d) PdL1 and (e) PdL4. (B).Solid state UV-Vis spectra of (a) PdL1 and (b) PdL1-Y.	122
4.9	(C) Solid state UV-Vis spectra of (a) PdL3 and (b) PdL3-Y. (D) Solid state UV-Vis spectra of (a) PdL4 and (b) PdL4-Y.	122
4.10	(E) Solid state UV-Vis spectra of (a) PdL5 and (b) PdL5-Y. (F) Solid state UV-Vis spectra of (a) PdL6 and (b) PdL6-Y.	122
5.1	Schematic representation of copper Schiff base complexes	131
5.2	Powder XRD patterns of (a) pure zeolite-Y (b) Cu- exchanged zeolite-Y, (c) CuL1-Y, (d) CuL3-Y, (e) CuL4-Y and (f) CuL6-Y	133
5.3	SEM images (A) CuL4-Y, before Soxhlet extraction and (B) CuL1-Y, (C) CuL4-Y after Soxhlet extraction.	134
5.4	(A) FTIR spectra of free state copper salen complexes (a) CuL1, (b) CuL3, (c) CuL4 and (d) CuL6. (B) FTIR spectra of encapsulated copper salen complexes (a) pure zeolite Y, (b) CuL1-Y, (c) CuL3-Y, (d) CuL4-Y and (e) CuL6-Y	135
5.5	XPS survey spectra of (a) CuL1 and (b) CuL1-Y	136
5.6	High resolution peak fitted XPS spectra of (A) Ni (2p) (B) C (1s), (C) N (1s) and (D) O(1s) for CuL1. (Black colored graphs are experimental data and red and blue colored peaks are peak fitted data)	137
5.7	(A). Solid state UV-Vis spectra of free state copper salen complexes (a) CuL1, (b) CuL3,(c) CuL4 and (d) CuL6 (B). The solid state UV-Vis spectra of encapsulated copper salen complexes(a) CuL1-Y, (b) CuL3-Y, (c) CuL4-Y and (d) CuL6-Y	140
5.8	Calibration curves of (A) styrene, (B) benzaldehyde and (C) styrene oxide	142

LIST OF FIGURES

5.9	Effect of substituent's on the geometry of complex around the metal and its effect on the transition state (T.S.) during the nucleophilic attack	147
6.1	Comparative molecular dimensions of cobalt Schiff-base complexes	152
6.2	SEM images of (A) pure zeolite Y, (B) Co-exchanged-Y, (C) CoL5-Y (before Soxhlet extraction) and (D) CoL5-Y (after Soxhlet extraction).	154
6.3	XRD pattern of (a) Pure zeolite-Y, (b) Co-exchanged zeolite -Y,(c)CoL1-Y, (d) CoL2-Y, (e) CoL3-Y, (f) CoL4-Y, (g) CoL5-Y and (h) CoL6-Y	155
6.4	(A)FTIR spectral data of cobalt Schiff-base complex in free state (a) CoL4, (b) CoL6, (c)CoL2, (d) CoL1, (e) CoL3 and (f) CoL5.(B) Enlarged view of in the range 1700- 800 cm ⁻¹ of cobalt Schiff-base complex in free state.	157
6.5	(A) FTIR spectral data of cobalt Schiff-base complex in free state (a) pure zeolite Y, (b)CoL6-Y, (c) CoL5-Y, (d) CoL3-Y, (e) CoL2-Y and (f) CoL1-Y and (g) CoL4-Y.(B) Enlarged view of in the range 1700- 900 cm ⁻¹ of cobalt Schiff-base complex in encapsulated state.	158
6.6	(Solid state UV-Visible spectra of (A) CoL1 (a), CoL1-Y (b). (B) CoL2 (a), CoL2-Y (b).(C) CoL3 (a), CoL3-Y(b). (D) CoL4 (a), CoL4-Y(b). (E) CoL5 (a), CoL5-Y (b). (F) CoL6 (a), CoL6-Y (b).	159
6.7	Magnetization vs. temperature plots in the range of 5K-300K of the complexes in free and encapsulated states (a) CoL1, (b) CoL4, (c) CoL1-Y (d) CoL6-Y and (e) CoL4-Y	162
6.8	Molecular structure of Rhodamine B (RhB)	163
6.9	The plot between percentage degradation and time taken by the different cobalt Schiff base complexes in free or encapsulated states under UV light irradiation (A) In absence of hydrogen peroxide (for 270 min) and (B) In presence of hydrogen (for 30 min) peroxide	164
7.1	Structure of NiL2 complex	170



This document was created with the Win2PDF "print to PDF" printer available at <http://www.win2pdf.com>

This version of Win2PDF 10 is for evaluation and non-commercial use only.

This page will not be added after purchasing Win2PDF.

<http://www.win2pdf.com/purchase/>