

# List of Figures

|     |  |    |
|-----|--|----|
| 2.1 | Evolution of Spin temperature . . . . .  | 13 |
| 2.2 | Evolution of the neutral fraction . . . . .  | 14 |
| 4.1 | Signal to noise ratio for the cross-correlation power spectrum for a single pointing radio observation of 400 hrs. We have considered a QSO density $30 \text{ deg}^{-2}$ . . . . .  | 49 |
| 4.2 | 68.3%,95.4% and 99.8% ellipse for 10000 hrs. observations for 25 pointings with each pointing of 400 hrs. The marginalized one dimensional probability distribution function (PDF) for $\Omega_m$ and $\Omega_\nu$ are also shown. . . . .   | 50 |
| 4.3 | Variation of $1-\sigma$ error of the parameters $\Omega_m$ and $\Omega_\nu$ with $N_{ant}$ the number of antennas in the radio-interferometric array. . . . .  | 52 |
| 4.4 | Variation of $1-\sigma$ error of the parameters $\Omega_m$ and $\Omega_\nu$ with $\bar{n}$ ( $\text{deg}^{-2}$ ). . . . .  | 53 |
| 4.5 | $\Omega_m$ contours in the $\bar{n}$ and time of observation plane. . . . .  | 54 |
| 4.6 | $\Omega_\nu$ contours in the $\bar{n}$ and time of observation plane. . . . .  | 54 |
| 5.1 | Shows the SNR for measuring the WDM cross power spectrum $P_{\mathcal{F}T}^{\text{WDM}}(k)$ in different $k$ -bins with observations of 200 hours each in $N_p = 25, 50$ and $100$ different fields-of-view. We have used a value of $m_{\text{WDM}} = 0.25 \text{ keV}$ for this analysis. The lower and upper horizontal lines in the figure correspond to $\text{SNR} = 5$ and $10$ respectively. . . . . | 65 |
| 5.2 | Shows the relative $1 - \sigma$ , $2 - \sigma$ and $3 - \sigma$ errors on the joint measurement of $\Omega_{\text{WDM}}$ and $\Omega_{\Lambda 0}$ with an observing time of 200 hours each in $N_p = 100$ different fields-of-view for $m_{\text{WDM}} = 0.25 \text{ keV}$ . . . . .   | 68 |

**LIST OF FIGURES**

---

|     |  |    |
|-----|--|----|
| 5.3 | Shows the $1 - \sigma$ errors (in red) on the measurement of the WDM cross power spectrum $P_{\mathcal{F}T}^{\text{WDM}}(k)$ in two different $k$ -bins with an observing time of 20000 hours distributed uniformly in $N_p = 100$ different fields-of-view. The upper, middle and lower panels in the figure show the results for the cases of $m_{\text{WDM}} = 0.20, 0.15$ and $0.10$ keV respectively. The WDM (blue line) and the matter (black line) cross power spectrum are plotted for the binned WDM (blue horizontal steps) and CDM (black horizontal steps) cross power spectrum respectively. . . . . | 70 |
| 5.4 | Figure showing the SNR for measuring the suppression in cross correlation power spectrum for a fiducial $m_{\text{WDM}} = 0.25$ Kev for a single pointing 21 cm observation for 400hrs, 2000hrs and 10000hrs respectively. . . . .   | 72 |
| 5.5 | Figures showing the 68.3%, 95.4% and 99.8% error contours for $\Omega_{\text{WDM}}$ and $\Omega_{\Lambda}$ . . . . .   | 73 |
| 6.1 | The growth rate of density perturbations for different CPL parametrizations. . . . .   | 77 |
| 6.2 | The growth rate of density perturbations for BA parametrizations .   | 78 |
| 6.3 | The dilation parameter $D_V$ for the different CPL parametrizations.   | 79 |
| 6.4 | The dilation parameter $D_V$ for the different BA parametrizations. .  | 79 |
| 6.5 | The CPL model ratio of $r_{\text{BAO}}$ with the LCDM model estimate . . .   | 80 |

# List of Tables

|     |  |    |
|-----|--|----|
| 4.1 | 1 $\sigma$ error for various number of pointings of the 21-cm observation for $\Omega_m$ with a total 10,000 hrs observation (distributed equally over the different pointing) of the 21 cm signal . . . . . | 51 |
| 4.2 | 1- $\sigma$ error for various number of pointings for $\Omega_\nu$ with a total 10,000 hrs observation (distributed equally over the different pointing) of the 21 cm signal. . . . .                        | 51 |
| 6.1 | Telescope specifications . . . . .   | 83 |
| 6.2 | The 1- $\sigma$ errors on $(w_0, w_a)$ from BAO imprint on the cross-correlation power spectrum. . . . .   | 84 |