

List of Figures

1.1	Melting Curve: Increase in fraction of open base pairs with temperature.	3
1.2	Force induced DNA melting: along the DNA axis & perpendicular to the DNA axis.	4
1.3	Mechanical stretching of DNA molecule through different experimental techniques.	5
2.1	Graphical presentation of PBD model. DNA double helix is simplified as a ladder. The filled circle represents the nucleotides, and the arrow shows that transverse motions are only considered.	14
2.2	(a) PBD model is viewed as a one-dimensional monotonic lattice model. (b) displacements of each nucleotides from their equilibrium positions.	15
2.3	Typical potential terms in a force field. Figures are taken from [1] .	22
3.1	The schematic representation of cations and DNA. The figure shows the probable distribution of cations around the two negative strands which are standing around 20\AA apart and the cations trying to nullify the repulsion between the two strands and possible distribution of excess cations that interact among themselves.	28
3.2	The specific heat as a function of temperature for the 12 base pair chain for the model parameters $k = 0.01 \text{ eV}/\text{\AA}^2$, $D_0 = 0.043 \text{ eV}$, $\lambda_1 = 0.01$, $\lambda_2 = -0.011$, $t = 0.01$, $\chi = 1.2$. The figure is plotted for $C = 0.5 \text{ M}$. The peak corresponds to the melting temperature. . . .	32
3.3	The temperature-salt phase diagrams showing the variation of T_m with salt concentration. The comparison of our results with the experimental results of [2]	33

LIST OF FIGURES

3.4 The phase diagram for different chain lengths in order to validate the results for infinite chain. The lengths are 36, 60, 156, 252, 276 bps respectively. 33

3.5 The critical force-salt phase diagram showing the variation of F_c with salt concentration. As there is no experimental data is available to compare, we show only the calculated points on the graph. . . . 35

4.1 (a) The schematic representation of the DNA molecule in a confined cell of conical shape. The distance x represents the distance between a pair and the confining wall while θ is the angle at which the cellular wall is suppose to confine the DNA. Here $z = 3.4 \times i$, the distance along x -axis. (b) The schematic representation of the DNA molecule in a confined cell of cylindrical geometry. The r is the distance of the confined wall from the DNA strand. So the diameter of the cylinder becomes $r + \text{DNA diameter}$ 39

4.2 (on left) The change in the melting temperature with the change in the angular separation between the confining wall and DNA molecule. The pore width $\delta = 0 \text{ \AA}$. Figure shows that beyond a certain angle, θ_c there is no effect of confining wall on the melting temperature, T_m of the molecule. (on right) The DNA is confined in a cylinder and the plot shows the change in melting temperature with radius of the cylinder. 40

4.3 The opening of the chain in different angular separation ($\theta = 5^\circ$ & 10°). When the angular separation is 5° , the effect lasts for ~ 25 base pairs while for the 10° angular separation, the effect lasts for ~ 20 base pairs at 380K. 42

4.4 The opening profile of the chain is shown through the 3D plots which shows a variation in $\langle y_j \rangle$ for the 120bp chain. Here we show the profile for two different angular separation, $\theta = 5^\circ$ and $\theta = 10^\circ$ (for pore width $\delta = 0 \text{ \AA}$) and for the cylindrical geometry $r = 20 \text{ \AA}$. The x -axis is site index while y -axis is temperature (in K) and z -axis denotes the average separation, $\langle y_j \rangle$ in \AA 43

4.5 How the effect of confinement changes with the linear size of the molecule is shown here. We consider the phage-lambda DNA molecule sequence of different lengths as 120, 180, 240 and 300. The pore diameter $\delta = 0 \text{ \AA}$ 44

LIST OF FIGURES

4.6	(On left) The change in the melting temperature (T_m) with the pore width (δ) for a fix value of angular separation ($\theta = 5^0$). (On right) The change in melting temperature (T_m) with the confinement angle (θ) for two values of the pore width (δ).	45
5.1	The schematic representation of the DNA chain confined in a cell. (a) Here, we schematically show a conically confined DNA molecule. The distance x represents the gap between a pair and the confining wall while θ is the angle at which the cellular wall is enclosing the DNA. Here $z = 3.4i$, the distance along x -axis. The parameter, δ , represents the pore width. (b) Here, we schematically show a cylindrically confined DNA molecule. The r is the distance of the confined wall from the DNA strand. So the radius of the cylinder becomes r +DNA radius.	51
5.2	The opening of the chain of 100 base pairs with temperature. The conical confined geometry makes the angle $\theta = 20^\circ$ and $\delta = 0\text{\AA}$.	52
5.3	The opening of the chain of 100 base pairs with temperature. Here we consider the separation between the cone vortex and the chain i.e $\delta = 0$, and $\theta = 20^\circ$.	53
5.4	The change in the transition temperature of the chain that is confined in a conical geometry. $\theta = 20^\circ$ for this graph. Here we consider two chains of different lengths, 100 and 200 base pairs.	55
5.5	The schematic of translocation process. We show the translocation of DNA through the three figures (a), (b) and (c). The fraction of pairs which stay inside/outside the cone varies as DNA comes out of the cone.	55
5.6	The plot shows the variation in the transition temperature, of DNA chain of 100 base pairs, with a fraction of chain outside the cell for different angular separations. Here the pore width, $\delta = 0\text{\AA}$.	56
5.7	The plot shows the variation in the transition temperature, of DNA chain of 100 base pairs, with a fraction of chain outside (inside) the cell for different angular separations.	58
5.8	The opening of the chain of 100 base pairs with temperature. The distance (r) is 20\AA and $\varphi = 50$. The confinement restriction starts from the base pair 50 ($\varphi = 50$), and hence inside the cone there are 50 base pairs while on the other side (out of the cylinder) there are 49 base pairs.	58

LIST OF FIGURES

5.9	The change in the transition temperature of the chain that is confined in a cylindrical geometry. The distance, $r = 10\text{\AA}$ is considered. We consider three chains of different lengths, 50, 100 and 200 base pairs.	59
5.10	DNA translocates through a pore of cylindrical geometry. The plot shows the transition temperature through different radial distance, r . Here the chain is of 100 base pairs.	60
5.11	The opening of the heterogeneous chain of 100 base pairs that is confined in a cylindrical geometry and the value of $r = 20\text{\AA}$ and $\varphi = 50$	61
5.12	The schematic representation of the confined geometry. For cylindrical confined geometry r represents the radius of the cylinder. . .	68
5.13	The rate of increasing of temperature with time.	68
5.14	RMSD calculation results are plotted for three cylindrical radius (r). The running average is shown for each RMSD fluctuations. . .	70
5.15	The location of the confinement. The radius of the confined cylinder for three systems are $r = 12.5\text{\AA}$	73
5.16	The RMSD calculation results are plotted for three different confined systems. The running average is shown for each RMSD fluctuations.	73
5.17	Nucleotide: The monomer of DNA polymer strand. The phosphate and carbonic base are bonded by covalent bond with the sugar. Image taken from <i>Wikipedia</i>	75
5.18	Nitrogenous bases: The two-ringed purines (Adenine and Guanine) and the single-ringed pyrimidines (Cytosine and Thymine). Thymine is unique to DNA. The diagram credit goes to 'The Medical Biochemistry Page', 'Lumen Learning'.	76
5.19	Basic structure of DNA, (single <i>strand</i>).	77
5.20	The secondary structure of DNA.	79
5.21	Schematic representation of DNA replication	81
5.22	Schematic representation of DNA replication	82
5.23	DNA Transcription	82

List of Tables

5.1	For $\varphi = 35, 50, 65$, we calculate the number of pairs that are open on the confined side (inside the cell) and the free side (outside the cell).	52
5.2	Counting Table of Open base pairs: $y_i > 1\text{\AA}$, is taken as open pair : we calculate the number of pairs that are open on the confined side (inside the cell) and free side (outside the cell) for $\delta = 0$ and $\theta = 20^\circ$	54
5.3	The average value of Intra-base pair parameters, Inter-base pair parameters and Base pair-axis parameters.	71
5.4	Comparison geometries of the most common DNA forms.	80

Nomenclature

Symbols

η	Crowder density
γ	Solution constant
κ	Single strand elasticity
λ	Solution constant
ϕ	Fraction of open pairs
ρ	Anharmonicity in stacking
θ	Fraction of intact pairs
a	Inverse of potential width
b	range of anharmonicity
C_v	Specific heat
D	Potential depth
f	Free energy
F_c	Critical force
H	Hamiltonian
k_β	Boltzmann constant
S	Entropy
T_m	Melting temperature
V_M	Morse potential

Z	Partition function
Z_c	Configurational part of the partition function
Z_p	Momentum part of the partition function

Acronyms

A	Adenine
AFM	Atomic force microscop
C	Cytocine
CEE	Constant extension ensemble
CFE	Constant force ensemble
DNA	Deoxyribonucleic Acid
dsDNA	Double stranded DNA
G	Guanine
MC	Monte Carlo
MD	Molecular Dynamics
mRNA	Messenger RNA
MT	Magnetic tweezer
OT	Optical tweezer
PBD	Peyrard Bishop Dauxois
RNA	Ribonucleic acid
SMFE	Single molecule force experiment
SMFS	Single molecule force spectroscopy
ssDNA	Single stranded DNA
T	Thymine
tRNA	Transfer RNA