

---



---

## LIST OF TABLES

---



---

No	Title	Page No
1.1	Optimization of reaction conditions for cross-dehydrogenative coupling reaction of heteroarenes	24
1.2	Substrate scope for the <b>XI</b> -catalyzed cross-dehydrogenative heterocoupling reaction for <b>44a-44zb</b>	26
1.3	Crystal data and structure refinement for complex <b>XI-XII</b>	42
2.2	Optimization of the reaction condition	63
2.3	Substrate scope for synthesis of 5-aryl-11 <i>H</i> -imidazo[1',2':1,2]-pyrido[3,4- <i>b</i> ]indoles ( <b>35</b> )	66
2.4	Substrate scope for synthesis of 5-aryl-7 <i>H</i> -imidazo[1',2':1,2]-pyrido[4,3- <i>b</i> ]indoles ( <b>38</b> )	67
2.5	Crystal data and structure refinement for <b>35ha</b>	84
2.6	Crystal data and structure refinement for <b>36aa</b>	85
3.3A.1	Optimization of reaction conditions	95
3.3A.2	Substrate scope for the synthesis of <b>13</b>	98
3.3A.3	Photophysical properties of the annulated products <b>13</b>	102
3.3A.4.5	Crystal data and structure refinement for <b>13ac</b>	118
3.3B.1	Optimization of reaction conditions	134
3.3B.2	Substrate scope for 2-arylindoles and maleimides	137
3.3B.3	Substrate scope for 2-heteroarylindoles	140
3.3B.4	Substrate scope for spirocyclization of 2,3-diphenylindoles	143
3.3B.5	Optical properties of <b>42</b> and <b>44</b>	147

3.3B.6	Crystal data and structure refinement for <b>42ab</b>	162
4.1	Optimization of reaction conditions for <i>ortho</i> -alkenylation	178
4.2	Alkenylation of 3-arylcoumarins	182
4.3	Regioselective halogenation of 3-arylcoumarins	183
4.4	Trifluoroethoxylation of 3-arylcoumarins	184
4.5	<i>Ortho</i> -Hydroxylation of 3-arylcoumarins	185
4.6	Synthesis of coumestan derivatives	185
4.7	Crystal data and structure refinement for <b>46aa</b>	216
4.8	Crystal data and structure refinement for <b>48ab</b>	217
4.9	Crystal data and structure refinement for <b>51a</b>	219

---

---

## LIST OF FIGURES

---

---

<b>Figure No.</b>	<b>Caption</b>	<b>Page No</b>
1.1	Traditional transition metal-catalyzed cross-coupling reactions	2
1.2	Transition metal-catalyzed C–H functionalization	2
1.3	Transition metal-catalyzed C–H functionalization with different coupling partners	3
1.4	Transition metal-catalyzed oxidative coupling	3
1.5	Flow chart presentation of cross-dehydrogenative coupling reactions	5
1.6	Classification of palladacycle	10
1.7	Examples of CE palladacycle	11
1.8	Examples of ECE palladacycle	11
1.9	Pincer ligands sites and their effects on the properties of the metal center	12
1.10	(A) $^1\text{H}$ NMR and (B) $^{13}\text{C}\{^1\text{H}\}$ NMR spectra of complex <b>XI</b> in $\text{CDCl}_3$ .	19
1.11	$^1\text{H}$ NMR and $^{13}\text{C}\{^1\text{H}\}$ NMR spectra of complex <b>XII</b> (complex trapped diethyl ether) in $\text{CDCl}_3$	20
1.12	$^1\text{H}$ NMR and $^{13}\text{C}\{^1\text{H}\}$ NMR spectra of complex <b>XIII</b> in $\text{CDCl}_3$	21
1.13	$^1\text{H}$ NMR and $^{13}\text{C}\{^1\text{H}\}$ NMR spectra of complex <b>XIV</b> in $\text{CDCl}_3$	22
1.14	a) Molecular structure of <b>XI</b> , b) Molecular structure of <b>XII</b> , c) Molecular structure of <b>XIII</b>	23
1.15	Catalytic life of <b>XI</b> under optimal conditions	28
2.1	Schematic representation of tandem reactions	49
2.2	Ring closure reaction <i>via</i> intramolecular oxidative C–H/ C–H coupling	50

2.3	<sup>1</sup> H NMR spectrum of 11-methyl-11 <i>H</i> -imidazo[1',2':1,2]-pyrido[3,4- <i>b</i> ]indol-5-yl)(phenyl)methanone ( <b>35aa</b> ) in CDCl <sub>3</sub>	59
2.4	<sup>13</sup> C{ <sup>1</sup> H} NMR spectrum of 11-methyl-11 <i>H</i> -imidazo[1',2':1,2]-pyrido[3,4- <i>b</i> ]indol-5-yl)(phenyl)methanone ( <b>35aa</b> ) in CDCl <sub>3</sub>	60
2.5	LC-HRMS of 11-methyl-11 <i>H</i> -imidazo[1',2':1,2]pyrido[3,4- <i>b</i> ]indol-5-yl)(phenyl)methanone ( <b>35aa</b> )	60
2.6	<sup>1</sup> H NMR spectrum of 2-(1 <i>H</i> -Imidazol-1-yl)-3-(1-methyl-1 <i>H</i> -indol-3-yl)-1-phenylprop-2-en-1-one ( <b>36aa</b> ) in CDCl <sub>3</sub>	61
2.7	<sup>13</sup> C{ <sup>1</sup> H} NMR spectrum of 2-(1 <i>H</i> -Imidazol-1-yl)-3-(1-methyl-1 <i>H</i> -indol-3-yl)-1-phenylprop-2-en-1-one ( <b>36aa</b> ) in CDCl <sub>3</sub>	61
2.8	LC-HRMS of 2-(1 <i>H</i> -Imidazol-1-yl)-3-(1-methyl-1 <i>H</i> -indol-3-yl)-1-phenylprop-2-en-1-one ( <b>36aa</b> )	62
2.9	Single crystal X-ray of 2-(1 <i>H</i> -Imidazol-1-yl)-3-(1-methyl-1 <i>H</i> -indol-3-yl)-1-phenylprop-2-en-1-one ( <b>36aa</b> )	62
2.10	Single-crystal X-ray of compound <b>35ha</b>	65
3.3A.1	Imidazo[1,2- <i>a</i> ]pyridine based drugs	89
3.3A.2	<sup>1</sup> H NMR spectra of 2-phenyl-1 <i>H</i> -benzo[ <i>e</i> ]pyrido[1',2':1,2]-imidazo[4,5- <i>g</i> ]isoindole-1,3(2 <i>H</i> )-dione ( <b>13aa</b> ) in CDCl <sub>3</sub>	96
3.3A.3	<sup>13</sup> C{ <sup>1</sup> H} NMR spectra of 2-phenyl-1 <i>H</i> -benzo[ <i>e</i> ]pyrido[1',2':1,2]-imidazo[4,5- <i>g</i> ]isoindole-1,3(2 <i>H</i> )-dione ( <b>13aa</b> ) in CDCl <sub>3</sub>	96
3.3A.4	HRMS spectrum of 2-phenyl-1 <i>H</i> -benzo[ <i>e</i> ]pyrido[1',2':1,2]-imidazo[4,5- <i>g</i> ]isoindole-1,3(2 <i>H</i> )-dione ( <b>13aa</b> )	97
3.3A.5	Single X-ray crystal structure of <b>13ac</b>	98

3.3A.6	UV-vis absorption (left) and normalized emission (right) spectra of a set of annulated compounds ( <b>13aa-ha</b> , <b>13ab</b> , <b>13ac</b> & <b>13ad</b> ) in CHCl <sub>3</sub> (1.0 × 10 <sup>-5</sup> M).	103
3.3A.7	Computed frontier molecular orbitals for <b>13aa</b> and <b>13ac</b>	104
3.3A.8	A) Absorption and B) normalized fluorescence spectra of <b>13aa</b> (1.0 × 10 <sup>-5</sup> M) in different solvents	105
3.3A.9	A) Absorption and B) fluorescence spectra of <b>13aa</b> (1.0 × 10 <sup>-5</sup> M in CHCl <sub>3</sub> ) in response to addition of TFA from 0 to 0.31 M.	105
3.3A.10	Selected illuminant compounds under visible light and UV irradiation at 365 nm	106
3.3B.1	Maleimide containing natural products and marine alkaloids	127
3.3B.2	<sup>1</sup> H and <sup>13</sup> C{ <sup>1</sup> H} NMR spectra of 2-methylbenzo[ <i>a</i> ]pyrrolo[3,4- <i>c</i> ]carbazole-1,3(2 <i>H</i> ,8 <i>H</i> )-dione ( <b>42aa</b> ) in DMSO- <i>d</i> <sub>6</sub>	135
3.3B.3	HRMS spectrum of 2-methylbenzo[ <i>a</i> ]pyrrolo[3,4- <i>c</i> ]carbazole-1,3(2 <i>H</i> ,8 <i>H</i> )-dione ( <b>42aa</b> )	136
3.3B.4	<sup>1</sup> H NMR spectrum of 5-methylpyrazolo[1',5':1,6]-pyrrolo[3',4':4,5]pyrido[2,3- <i>b</i> ]indole-4,6(5 <i>H</i> ,11 <i>H</i> )-dione ( <b>44aa</b> ) in DMSO- <i>d</i> <sub>6</sub>	139
3.3B.5	<sup>13</sup> C{ <sup>1</sup> H} NMR spectrum of 5-methylpyrazolo[1',5':1,6]-pyrrolo[3',4':4,5]pyrido[2,3- <i>b</i> ]indole-4,6(5 <i>H</i> ,11 <i>H</i> )-dione ( <b>44aa</b> ) in DMSO- <i>d</i> <sub>6</sub>	139
3.3B.6	HRMS spectrum of 5-methylpyrazolo[1',5':1,6]-pyrrolo[3',4':4,5]pyrido[2,3- <i>b</i> ]indole-4,6(5 <i>H</i> ,11 <i>H</i> )-dione ( <b>44aa</b> )	140
3.3B.7	<sup>1</sup> H NMR spectra of 1'-methyl-11-phenylspiro-[isoindolo[2,1- <i>a</i> ]indole-6,3'-pyrrolidine]-2',5'-dione ( <b>46aa</b> ) in DMSO- <i>d</i> <sub>6</sub>	141
3.3B.8	<sup>13</sup> C{ <sup>1</sup> H} NMR spectra of 1'-methyl-11-phenylspiro-[isoindolo[2,1- <i>a</i> ]indole-6,3'-pyrrolidine]-2',5'-dione ( <b>46aa</b> ) in DMSO- <i>d</i> <sub>6</sub>	142

3.3B.9	HRMS spectrum of 1'-methyl-11-phenylspiro[isoindolo[2,1- <i>a</i> ]indole-6,3'-pyrrolidine]-2',5'-dione ( <b>46aa</b> )	142
3.3B.10	Normalized a) UV-visible and b) emission spectra of <b>14a</b> , <b>42aa</b> , <b>44aa</b> and representative c) UV-visible and d) emission spectra of <b>42</b> and <b>44</b> in DMSO	146
3.3B.11	ORTEP diagram of <b>42ab</b> (CCDC No 2024848). The thermal ellipsoids are drawn at 50% probability level	162
4.1	C–H functionalization through cyclometallation of directing groups	168
4.2	Representative examples of strong and weakly directing groups	169
4.3	<sup>1</sup> H and <sup>13</sup> C{ <sup>1</sup> H} NMR spectra of methyl ( <i>E</i> )-3-(2-(2-oxo-2 <i>H</i> -chromen-3-yl)phenyl)acrylate ( <b>46aa</b> ) in CDCl <sub>3</sub>	180
4.4	HRMS spectrum of methyl ( <i>E</i> )-3-(2-(2-oxo-2 <i>H</i> -chromen-3-yl)phenyl)acrylate ( <b>46aa</b> )	181
4.5	ORTEP diagram of compound <b>46aa</b> , the thermal ellipsoids are drawn to the 50% probability level	215
4.6	ORTEP diagram of compound <b>48ab</b> , the thermal ellipsoids are drawn to the 50% probability level	217
4.7	ORTEP diagram of compound <b>51a</b> , the thermal ellipsoids are drawn to the 50% probability level	218
5.1	Flow chart presentation of cross-dehydrogenative coupling reactions	225
5.2	Transition metal-catalyzed synthesis of fused heterocycles	231

## LIST OF ABBREVIATIONS

Abbreviation	Description
AcOH	Acetic acid
acac	Acetylacetone
ADA	1-Adamantancarboxylic acid
Aq	Aqueous
Ar	Aryl
BHT	Butylated hydroxy toluene
<i>t</i> -BuOK	Potassium <i>tert</i> -butoxide
<i>t</i> -BuONO	<i>tert</i> -Butyl nitrite
Cat.	Catalytic
CMD	Concerted metalation deprotonation
<sup>13</sup> C	Carbon-13
CH <sub>3</sub> CN	Acetonitrile
CDCl <sub>3</sub>	Deuterated chloroform
<i>m</i> -CPBA	<i>meta</i> -Chloroperoxybenzoic acid
CDC	Cross-dehydrogenative coupling
CAN	Ceric ammonium nitrate
Cp*	1,2,3,4,5-Pentamethylcyclopenta-1,3-diene
<i>p</i> -Cymene	1-Methyl-4-(propan-2-yl)benzene
DMAP	4-Dimethylaminopyridine
d	Doublet
DBU	1,8-Diazabicyclo[5.4.0]undec-7-ene

dd	Doublet of doublet
DCE	1,2-Dichloroethane
DCM	Dichloromethane
DMA	<i>N,N</i> -Dimethylacetamide
DME	Dimethoxyethane
DMF	<i>N,N</i> -Dimethylformamide
DMSO- <i>d</i> <sub>6</sub>	Deuterated dimethylsulfoxide
DMSO	Dimethylsulfoxide
DMAc	Dimethylacetamide
DMAP	4-Dimethylaminopyridine
DDQ	2,3-Dichloro-5,6-dicyano-1,4-benzoquinone
DGs	Directing groups
dppf	1,1'-Bis(diphenylphosphino)ferrocene
EDC.HCl	1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide
EI	Electron ionization
EtOAc	Ethyl acetate
equiv.	Equivalent
ESI-MS	Electrospray ionization mass spectrometry
EtOH	Ethanol
FID	Free induction decay
FT-IR	Fourier transform infrared
g	Gram
h	Hours

HFIP	1,1,1,3,3,3-Hexafluoro-2-propanol
HRMS	High resolution mass spectra
Hz	Hertz
HOMO	Highest occupied molecular orbital
IR	Infra-red
IPA	Isopropanol
IBD	Iodobenzene diacetate
IOC	Intramolecular oxidative coupling
IP	Imidazo[1,2- <i>a</i> ]pyridines
<i>J</i>	Coupling constant
KOAc	Potassium acetate
KIE	Kinetic isotope effect
Lit.	Literature
LUMO	Lowest unoccupied molecular orbital
mp	Melting point
<i>m</i>	Multiplet
mg	Milligram
MHz	Mega hertz
min	Minutes
mmol	Millimole
MW	Microwave
nm	Nanometer
N <sub>2</sub>	Nitrogen
DIPEA	N,N-Diisopropylethylamine

Nu	Nucleophile
NMR	Nuclear magnetic resonance
NBS	<i>N</i> -Bromosuccinimide
NCS	<i>N</i> -Chlorosuccinimide
NIS	<i>N</i> -Iodoosuccinimide
NMP	<i>N</i> -Methyl-2-pyrrolidone
O <sub>2</sub>	Oxygen
OTf	Trifluoromethanesulfonate
OLEDs	Organic light-emitting diode
PEG400	Polyethylene glycol 400
PIDA	Phenyliodine(III) diacetate
1,10-Phen	1,10-Phenanthroline
PivOH	Pivalic acid
PhCl	Chlorobenzene
ppm	Parts per million
r.t.	Room temperature
s	Singlet
SET	Single electron transfer
S <sub>E</sub> Ar	Substitution electrophilic aromatic
TFA	Trifluoroacetic acid
TFE	2,2,2-Trifluoroethanol
TEA	Triethyl amine
t	Triplet
TMDEA	Tetramethylethylenediamine

TBAB	Tetrabutylammonium bromide
TBHP	<i>tert</i> -Butyl hydroperoxide
THF	Tetrahydrofuran
TEMPO	2,2,6,6-Tetramethyl-1-piperidinyloxy
<i>p</i> -TSA or <i>p</i> -TsOH	<i>p</i> -Toluenesulfonic acid
qTOF	Quadrupole time-of-flight
TLC	Thin layer chromatography
TfOH	Trifluoromethanesulfonic acid
TFAA	Trifluoroacetic anhydride
TMS	Tetramethylsilane
TMSCl	Trimethylsilyl chloride
TD-DFT	Time dependent density functional theory
UV	Ultraviolet-visible