

9. References

- Abadir PM, Walston JD, & Carey RM (2012). Subcellular characteristics of functional intracellular renin-angiotensin systems. *Peptides* 38: 437-445.
- Abdellatif SA, Galal AA, Farouk SM, & Abdel-Daim MM (2017). Ameliorative effect of parsley oil on cisplatin-induced hepato-cardiotoxicity: a biochemical, histopathological, and immunohistochemical study. *Biomedicine & Pharmacotherapy* 86: 482-491.
- Adachi N, Lei B, Deshpande G, Seyfried FJ, Shimizu I, Nagaro T, *et al.* (2001). Uraemia suppresses central dopaminergic metabolism and impairs motor activity in rats. *Intensive Care Medicine* 27: 1655-1660.
- Adachi T, Sugiyama N, Gondai T, Yagita H, & Yokoyama T (2013). Blockade of death ligand TRAIL inhibits renal ischemia reperfusion injury. *Acta Histochemica et Cytochemica* 46: 161-170.
- Ahmed HA, Ishrat T, Pillai B, Bunting KM, Vazdarjanova A, Waller JL, *et al.* (2019). Angiotensin receptor (AT2R) agonist C21 prevents cognitive decline after permanent stroke in aged animals—A randomized double-blind pre-clinical study. *Behavioural Brain Research* 359: 560-569.
- Alabdan N, Gosmanova EO, Tran NQT, Oliphant CS, Pan H, Broyles JE, *et al.* (2016). Risk Factors for Acute Kidney Injury in Patients Continued on Renin-Angiotensin System Blockers During Hospitalization. *The American Journal of the Medical Sciences* 353: 172-177.
- Albiston AL, Diwakarla S, Fernando RN, Mountford SJ, Yeatman HR, Morgan B, *et al.* (2011). Identification and development of specific inhibitors for insulin-regulated aminopeptidase as a new class of cognitive enhancers. *British Journal of Pharmacology* 164: 37-47.
- Alge JL, Karakala N, Neely BA, Janech MG, Tumlin JA, Chawla LS, *et al.* (2013). Urinary angiotensinogen and risk of severe AKI. *Clinical Journal of the American Society of Nephrology* 8: 184-193.

- Allikmets K (2007). Aliskiren--an orally active renin inhibitor. Review of pharmacology, pharmacodynamics, kinetics, and clinical potential in the treatment of hypertension. *Vascular Health and Risk Management* 3: 809-815.
- Alobaidi R, Basu RK, Goldstein SL, & Bagshaw SM Sepsis-associated acute kidney injury. *Seminar in Nephrology* 35:2-11.
- Amini N, Sarkaki A, Dianat M, Mard SA, Ahangarpour A, & Badavi M (2019). Protective effects of naringin and trimetazidine on remote effect of acute renal injury on oxidative stress and myocardial injury through Nrf-2 regulation. *Pharmacological Reports* 71: 1059-1066.
- Amura CR, Renner B, Lyubchenko T, Faubel S, Simonian PL, & Thurman JM (2012). Complement activation and toll-like receptor-2 signaling contribute to cytokine production after renal ischemia/reperfusion. *Molecular Immunology* 52: 249-257.
- Anacker C, & Hen R (2017). Adult hippocampal neurogenesis and cognitive flexibility—linking memory and mood. *Nature Reviews Neuroscience* 18: 335-346.
- Andres-Hernando A, Dursun B, Altmann C, Ahuja N, He Z, Bhargava R, *et al.* (2012). Cytokine production increases and cytokine clearance decreases in mice with bilateral nephrectomy. *Nephrology Dialysis Transplantation* 27: 4339-4347.
- Anker SD, & Coats AJ (2002). How to RECOVER from RENAISSANCE? The significance of the results of RECOVER, RENAISSANCE, RENEWAL and ATTACH. *International Journal of Cardiology* 86: 123-130.
- Arany I, Herbert J, Herbert Z, & Safirstein RL (2008). Restoration of CREB function ameliorates cisplatin cytotoxicity in renal tubular cells. *American Journal of Physiology-Renal Physiology* 294: F577-F581.
- Arroyo D, Melero R, Panizo N, Goicoechea M, Rodríguez-Benítez P, Vinuesa SG, *et al.* (2011). Metformin-associated acute kidney injury and lactic acidosis. *International Journal of Nephrology* 2011: 1-5.

- Ascon M, Ascon DB, Liu M, Cheadle C, Sarkar C, Racusen L, *et al.* (2009). Renal ischemia-reperfusion leads to long term infiltration of activated and effector-memory T lymphocytes. *Kidney International* 75: 526-535.
- Awad AS, Kamel R, & Sherief MAE (2011). Effect of thymoquinone on hepatorenal dysfunction and alteration of CYP3A1 and spermidine/spermine N-1-acetyl-transferase gene expression induced by renal ischaemia–reperfusion in rats. *Journal of Pharmacy and Pharmacology* 63: 1037-1042.
- Azizi M, & Menard J (2013). Renin inhibitors and cardiovascular and renal protection: an endless quest? *Cardiovascular Drugs and Therapy* 27: 145-153.
- Ba Aqeel SH, Sanchez A, & Batlle D (2017). Angiotensinogen as a biomarker of acute kidney injury. *Clinical Kidney Journal* 10: 759-768.
- Bader M (2013). ACE2, angiotensin-(1-7), and Mas: the other side of the coin. *Pflugers Archiv : European Journal of Physiology* 465: 79-85.
- Bader M, Santos RA, Unger T, & Steckelings UM (2012). New therapeutic pathways in the RAS. *Journal of the Renin-angiotensin-aldosterone System* 13: 505-508.
- Bae EH, Fang F, Williams VR, Konvalinka A, Zhou X, Patel VB, *et al.* (2017). Murine recombinant angiotensin-converting enzyme 2 attenuates kidney injury in experimental Alport syndrome. *Kidney International* 91: 1347-1361.
- Bagshaw SM, Hoste EA, Braam B, Briguori C, Kellum JA, McCullough PA, *et al.* (2013). Cardiorenal syndrome type 3: pathophysiologic and epidemiologic considerations. In ADQI Consensus on AKI Biomarkers and Cardiorenal Syndromes. Karger Publishers, pp 137-157.
- Bakker PJ, Scantleberry AM, Butter LM, Claessen N, Teske GJ, van der Poll T, *et al.* (2015). TLR9 mediates remote liver injury following severe renal ischemia reperfusion. *PloS one* 10: e0137511.
- Bannister AJ, & Kouzarides T (2011). Regulation of chromatin by histone modifications. *Cell research* 21: 381-395.

- Barrera Chimal J, Perez Villalva R, Rodriguez-Romo R, Reyna J, Uribe N, Gamba G, *et al.* (2013). Spironolactone prevents chronic kidney disease caused by ischemic acute kidney injury. *Kidney International* 83: 93-103.
- Barroso LC, Silveira KD, Lima CX, Borges V, Bader M, Rachid M, *et al.* (2012). Renoprotective effects of AVE0991, a nonpeptide Mas receptor agonist, in experimental acute renal injury. *International Journal of Hypertension* 2012: 1-8.
- Basile DP, Bonventre JV, Mehta R, Nangaku M, Unwin R, Rosner MH, *et al.* (2016). Progression after AKI: Understanding Maladaptive Repair Processes to Predict and Identify Therapeutic Treatments. *Journal of the American Society of Nephrology* 27: 687-697.
- Basile DP, Leonard EC, Beal AG, Schleuter D, & Friedrich J (2012). Persistent oxidative stress following renal ischemia-reperfusion injury increases ANG II hemodynamic and fibrotic activity. *American Journal of Physiology-Renal Physiology* 302: 1494-1502.
- Basso N, & Terragno NA (2001). History about the discovery of the renin-angiotensin system. *Hypertension* 38: 1246-1249.
- Bavishi C, Bangalore S, & Messerli FH (2016). Renin angiotensin aldosterone system inhibitors in hypertension: is there evidence for benefit independent of blood pressure reduction? *Progress in Cardiovascular Diseases* 59: 253-261.
- Bayrak O, Uz E, Bayrak R, Turgut F, Atmaca AF, Sahin S, *et al.* (2008). Curcumin protects against ischemia/reperfusion injury in rat kidneys. *World Journal of Urology* 26: 285-291.
- Becker LB (2004). New concepts in reactive oxygen species and cardiovascular reperfusion physiology. *Cardiovascular Research* 61: 461-470.
- Beker BM, Corleto MG, Fieiras C, & Musso CG (2018). Novel acute kidney injury biomarkers: their characteristics, utility and concerns. *International Urology and Nephrology* 50: 705-713.
- Bellomo R, Kellum JA, & Ronco C (2012). Acute kidney injury. *The Lancet* 380: 756-766.

- Bennion DM, Haltigan EA, Irwin AJ, Donnangelo LL, Regenhardt RW, Pioquinto DJ, *et al.* (2015). Activation of the Neuroprotective Angiotensin-Converting Enzyme 2 in Rat Ischemic Stroke. *Hypertension (Dallas, Tex : 1979)* 66: 141-148.
- Bennion DM, Isenberg JD, Harmel AT, DeMars K, Dang AN, Jones CH, *et al.* (2017). Post-stroke angiotensin II type 2 receptor activation provides long-term neuroprotection in aged rats. *PloS one* 12: e0180738.
- Bernstein KE, Sayeski PP, Doan T, Farmer P, & Ali MS (2001). Signal transduction pathways of angiotensin II in the kidney. In *The Renin-angiotensin System and Progression of Renal Diseases*. Karger Publishers, pp 16-33.
- Berry C, Touyz R, Dominiczak AF, Webb RC, & Johns DG (2001). Angiotensin receptors: signaling, vascular pathophysiology, and interactions with ceramide. *American Journal of Physiology Heart and Circulatory Physiology* 281: 2337-2365.
- Bertagnolli M, Casali KR, De Sousa FB, Rigatto K, Becker L, Santos SH, *et al.* (2014). An orally active angiotensin-(1-7) inclusion compound and exercise training produce similar cardiovascular effects in spontaneously hypertensive rats. *Peptides* 51: 65-73.
- Bhaumik SR, Smith E, & Shilatifard A (2007). Covalent modifications of histones during development and disease pathogenesis. *Nature Structural & Molecular Biology* 14: 1008-1016.
- Bigagli E, & Lodovici M (2019). Circulating Oxidative Stress Biomarkers in Clinical Studies on Type 2 Diabetes and Its Complications. *Oxidative Medicine and Cellular Longevity* 2019: 1-18.
- Bihorac A, Chawla LS, Shaw AD, Al-Khafaji A, Davison DL, Demuth GE, *et al.* (2014). Validation of cell-cycle arrest biomarkers for acute kidney injury using clinical adjudication. *American Journal of Respiratory and Critical Care Medicine* 189: 932-939.
- Bird A (2002). DNA methylation patterns and epigenetic memory. *Genes & Development* 16: 6-21.

- Bird L, & Walker D (2015). Pathophysiology of acute kidney injury. Companion Animal 20: 142-147.
- Blanco VE, Hernandorena CV, Scibona P, Beloso W, & Musso CG (2019). Acute Kidney Injury Pharmacokinetic Changes and Its Impact on Drug Prescription. Healthcare 7: 10.
- Bockenkamp B, & Vyas H (2003). Understanding and managing acute fluid and electrolyte disturbances. Current Paediatrics 13: 520-528.
- Bomsztyk K, Flanigan S, Mar D, Mikula M, Johnson A, Zager R, *et al.* (2013). Synchronous recruitment of epigenetic modifiers to endotoxin synergistically activated Tnf- α gene in acute kidney injury. PloS one 8: e70322-e70322.
- Bonventre JV, & Yang L (2011). Cellular pathophysiology of ischemic acute kidney injury. The Journal of Clinical Investigation 121: 4210-4221.
- Bordi M, Berg MJ, Mohan PS, Peterhoff CM, Alldred MJ, Che S, *et al.* (2016). Autophagy flux in CA1 neurons of Alzheimer hippocampus: Increased induction overburdens failing lysosomes to propel neuritic dystrophy. Autophagy 12: 2467-2483.
- Bouchard J, Soroko SB, Chertow GM, Himmelfarb J, Ikizler TA, Paganini EP, *et al.* (2009). Fluid accumulation, survival and recovery of kidney function in critically ill patients with acute kidney injury. Kidney International 76: 422-427.
- Boyne AF, & Ellman GL (1972). A methodology for analysis of tissue sulfhydryl components. Analytical Biochemistry 46: 639-653.
- Brasacchio D, Okabe J, Tikellis C, Balcerzyk A, George P, Baker EK, *et al.* (2009). Hyperglycemia induces a dynamic cooperativity of histone methylase and demethylase enzymes associated with gene-activating epigenetic marks that coexist on the lysine tail. Diabetes 58: 1229-1236.
- Braszko JJ, Walesiuk A, & Wielgat P (2006). Cognitive effects attributed to angiotensin II may result from its conversion to angiotensin IV. Journal of the Renin-Angiotensin-Aldosterone System 7: 168-174.

- Brilli LL, Swanhart LM, de Caestecker MP, & Hukriede NA (2013). HDAC inhibitors in kidney development and disease. *Pediatric Nephrology* 28: 1909-1921.
- Brochner AC, Dagnaes-Hansen F, Hojberg-Holm J, & Toft P (2014). The inflammatory response in blood and in remote organs following acute kidney injury. *APMIS : Acta Pathologica, Microbiologica, et Immunologica Scandinavica* 122: 399-404.
- Bruce E, Shenoy V, Rathinasabapathy A, Espejo A, Horowitz A, Oswalt A, *et al.* (2015). Selective activation of angiotensin AT2 receptors attenuates progression of pulmonary hypertension and inhibits cardiopulmonary fibrosis. *British Journal of Pharmacology* 172: 2219-2231.
- Bucsics T, & Krones E (2017). Renal dysfunction in cirrhosis: acute kidney injury and the hepatorenal syndrome. *Gastroenterology Report* 5: 127-137.
- Burn D, & Bates D (1998). Neurology and the kidney. *Journal of Neurology, Neurosurgery & Psychiatry* 65: 810-821.
- Campbell DJ (2014). Clinical relevance of local renin angiotensin systems. *Frontiers in Endocrinology* 5: 1-5.
- Cao Q, Harris DC, & Wang Y (2015a). Macrophages in kidney injury, inflammation, and fibrosis. *Physiology* 30: 183-194.
- Cao R, Tsukada Y-i, & Zhang Y (2005). Role of Bmi-1 and Ring1A in H2A ubiquitylation and Hox gene silencing. *Molecular Cell* 20: 845-854.
- Cao W, Li A, Li J, Wu C, Cui S, Zhou Z, *et al.* (2017). Reno-cerebral reflex activates the renin-angiotensin system, promoting oxidative stress and renal damage after ischemia-reperfusion injury. *Antioxidants & Redox Signaling* 27: 415-432.
- Cao W, Li A, Wang L, Zhou Z, Su Z, Bin W, *et al.* (2015b). A salt-induced reno-cerebral reflex activates renin-angiotensin systems and promotes CKD progression. *Journal of the American Society of Nephrology* 26: 1619-1633.

- Cao X, Song LN, Zhang YC, Li Q, Shi TT, Yang FY, *et al.* (2019). Angiotensin-converting enzyme 2 inhibits endoplasmic reticulum stress-associated pathway to preserve nonalcoholic fatty liver disease. *Diabetes/Metabolism Research and Reviews* 35: e3123.
- Cardenas A (2013). Defining renal failure in cirrhosis-Acute kidney injury classification or traditional criteria? *Annals of Hepatology* 12: 984-985.
- Caretti G, Di Padova M, Micales B, Lyons GE, & Sartorelli V (2004). The Polycomb Ezh2 methyltransferase regulates muscle gene expression and skeletal muscle differentiation. *Genes & Development* 18: 2627-2638.
- Carey RM (2005). Cardiovascular and renal regulation by the angiotensin type 2 receptor: the AT2 receptor comes of age. *Hypertension* 45: 840-844.
- Carey RM (2017a). AT2 receptors: potential therapeutic targets for hypertension. *American Journal of Hypertension* 30: 339-347.
- Carey RM (2017b). Update on angiotensin AT2 receptors. *Current Opinion in Nephrology and Hypertension* 26: 91-96.
- Castellano G, Franzin R, Stasi A, Divella C, Sallustio F, Pontrelli P, *et al.* (2018). Complement Activation During Ischemia/Reperfusion Injury Induces Pericyte-to-Myofibroblast Transdifferentiation Regulating Peritubular Capillary Lumen Reduction Through pERK Signaling. *Frontiers in Immunology* 9: 1002.
- Castoldi G, di Gioia CR, Bombardi C, Maestroni S, Carletti R, Steckelings UM, *et al.* (2014). Prevention of diabetic nephropathy by compound 21, selective agonist of angiotensin type 2 receptors, in Zucker diabetic fatty rats. *American Journal of Physiology-Renal Physiology* 307: 1123-1131.
- Challa A, Chan W, Krieg Jr RJ, Thabet MA, Liu F, Hintz RL, *et al.* (1993). Effect of metabolic acidosis on the expression of insulin-like growth factor and growth hormone receptor. *Kidney International* 44: 1224-1227.

- Chawla LS, Eggers PW, Star RA, & Kimmel PL (2014). Acute kidney injury and chronic kidney disease as interconnected syndromes. *The New England Journal of Medicine* 371: 58-66.
- Chen H, Wan D, Wang L, Peng A, Xiao H, Petersen RB, *et al.* (2015). Apelin protects against acute renal injury by inhibiting TGF- β 1. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease* 1852: 1278-1287.
- Chen J, Guo Y, Zeng W, Huang L, Pang Q, Nie L, *et al.* (2014a). ER stress triggers MCP-1 expression through SET7/9-induced histone methylation in the kidneys of db/db mice. *American Journal of Physiology-Renal Physiology* 306: F916-F925.
- Chen J, Zhao Y, Chen S, Wang J, Xiao X, Ma X, *et al.* (2014b). Neuronal over-expression of ACE2 protects brain from ischemia-induced damage. *Neuropharmacology* 79: 550-558.
- Chen R, Kang R, Fan X, & Tang D (2014c). Release and activity of histone in diseases. *Cell death & disease* 5: e1370-e1370.
- Chen S, Bellew C, Yao X, Stefkova J, Dipp S, Saifudeen Z, *et al.* (2011). Histone deacetylase (HDAC) activity is critical for embryonic kidney gene expression, growth, and differentiation. *Journal of Biological Chemistry* 286: 32775-32789.
- Chen TH, Yang YC, Wang JC, & Wang JJ Curcumin Treatment Protects Against Renal Ischemia and Reperfusion Injury-Induced Cardiac Dysfunction and Myocardial Injury. 45 (10):3546-3549.
- Chen WL, Sun HP, Li DD, Wang ZH, You QD, & Guo XK (2017). G9a-an appealing antineoplastic target. *Current Cancer Drug Targets* 17: 555-568.
- Chen W, Tang D, Xu Y, Zou Y, Sui W, Dai Y, *et al.* (2018). Comprehensive analysis of lysine crotonylation in proteome of maintenance hemodialysis patients. *Medicine* 97:37(e12035).
- Cherblanc FL, Chapman KL, Reid J, Borg AJ, Sundriyal S, Alcazar-Fuoli L, *et al.* (2013). On the histone lysine methyltransferase activity of fungal metabolite chaetocin. *Journal of Medicinal Chemistry* 56: 8616-8625.

- Chiang WC, Chien CT, Lin WW, Lin SL, Chen YM, Lai CF, *et al.* (2006). Early activation of bradykinin B2 receptor aggravates reactive oxygen species generation and renal damage in ischemia/reperfusion injury. *Free Radical Biology and Medicine* 41: 1304-1314.
- Choi DE, Jeong JY, Lim BJ, Chang YK, Na KR, Shin YT, *et al.* (2011). Aliskiren ameliorates renal inflammation and fibrosis induced by unilateral ureteral obstruction in mice. *The Journal of Urology* 186: 694-701.
- Choi DE, Jeong JY, Lim BJ, Chung S, Chang YK, Lee SJ, *et al.* (2009). Pretreatment of sildenafil attenuates ischemia-reperfusion renal injury in rats. *American Journal of Physiology-Renal Physiology* 297: F362-F370.
- Chouchani ET, Pell VR, James AM, Work LM, Saeb-Parsy K, Frezza C, *et al.* (2016). A unifying mechanism for mitochondrial superoxide production during ischemia-reperfusion injury. *Cell Metabolism* 23: 254-263.
- Choudhary C, Kumar C, Gnad F, Nielsen ML, Rehman M, Walther TC, *et al.* (2009). Lysine acetylation targets protein complexes and co-regulates major cellular functions. *Science* 325: 834-840.
- Chow BS, Koulis C, Krishnaswamy P, Steckelings UM, Unger T, Cooper ME, *et al.* (2016). The angiotensin II type 2 receptor agonist Compound 21 is protective in experimental diabetes-associated atherosclerosis. *Diabetologia* 59: 1778-1790.
- Chrysostomou A, Pedagogos E, MacGregor L, & Becker GJ (2006). Double-blind, placebo-controlled study on the effect of the aldosterone receptor antagonist spironolactone in patients who have persistent proteinuria and are on long-term angiotensin-converting enzyme inhibitor therapy, with or without an angiotensin II receptor blocker. *Clinical Journal of American Society of Nephrology* 1: 256-262.
- Classics Lowry O, Rosebrough N, Farr A, & Randall R (1951). Protein measurement with the Folin phenol reagent. *Journal of biological Chemistry* 193: 265-275.

- Collaboration NRF (2016). Worldwide trends in diabetes since 1980: a pooled analysis of 751 population-based studies with 4.4 million participants. *Lancet* (London, England) 387: 1513-1530.
- Coorevits P, Sundgren M, Klein GO, Bahr A, Claerhout B, Daniel C, *et al.* (2013). Electronic health records: new opportunities for clinical research. *Journal of Internal Medicine* 274: 547-560.
- Cortes AL, Gonsalez SR, Rioja LS, Oliveira SS, Santos AL, Prieto MC, *et al.* (2018). Protective outcomes of low-dose doxycycline on renal function of Wistar rats subjected to acute ischemia/reperfusion injury. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease* 1864: 102-114.
- Corvol P, Eyries M, & Soubrier F (2004). Peptidyl-dipeptidase A/angiotensin I-converting enzyme. In *Handbook of Proteolytic Enzymes* (Second Edition), Vol 1. Elsevier, pp 332-346.
- Cosentino CC, Skrypnyk NI, Brilli LL, Chiba T, Novitskaya T, Woods C, *et al.* (2013). Histone deacetylase inhibitor enhances recovery after AKI. *Journal of the American Society of Nephrology* 24: 943-953.
- Costalanga EC, Silva FMO, & Noronha IL (2016). Valproic Acid Prevents Renal Dysfunction and Inflammation in the Ischemia-Reperfusion Injury Model. *Biomed Research International* 2016: 1-10.
- Crackower MA, Sarao R, Oudit GY, Yagil C, Kozieradzki I, Scanga SE, *et al.* (2002). Angiotensin-converting enzyme 2 is an essential regulator of heart function. *Nature* 417: 822-828.
- Craddock GN (1976). Species differences in response to renal ischemia. *Archives of Surgery* 111: 582-584.
- da Fonseca Magalhães PA, de Brito TS, Freire RS, da Silva MTB, dos Santos AA, Vale ML, *et al.* (2016). Metabolic acidosis aggravates experimental acute kidney injury. *Life Sciences* 146: 58-65.

- da Silveira KD, Bosco KSP, Diniz LR, Carmona AK, Cassali GD, Bruna-Romero O, *et al.* (2010). ACE2–angiotensin-(1–7)–Mas axis in renal ischaemia/reperfusion injury in rats. *Clinical Science* 119: 385-394.
- Dai S-Y, Peng W, Zhang Y-P, Li J-D, Shen Y, & Sun X-F (2015). Brain endogenous angiotensin II receptor type 2 (AT2-R) protects against DOCA/salt-induced hypertension in female rats. *Journal of Neuroinflammation* 12: 1-8.
- Dai S-Y, Zhang Y-P, Peng W, Shen Y, & He J-J (2016). Central infusion of angiotensin II type 2 receptor agonist compound 21 attenuates DOCA/NaCl-induced hypertension in female rats. *Oxidative Medicine and Cellular Longevity* 2016: 1-9.
- Danelli L, Madjene LC, Madera-Salcedo I, Gautier G, Pacreau E, Mkaddem SB, *et al.* (2017). Early phase mast cell activation determines the chronic outcome of renal ischemia–reperfusion injury. *The Journal of Immunology* 198: 2374-2382.
- Danobeitia JS, Ziemelis M, Ma X, Zitur LJ, Zens T, Chlebeck PJ, *et al.* (2017). Complement inhibition attenuates acute kidney injury after ischemia-reperfusion and limits progression to renal fibrosis in mice. *PloS one* 12 e0183701.
- De Gasparo M, Catt K, Inagami T, Wright J, & Unger T (2000). International union of pharmacology. XXIII. The angiotensin II receptors. *Pharmacological Reviews* 52: 415-472.
- De Luca Jr LA, Menani JV, & Johnson AK (2013) *Neurobiology of Body Fluid Homeostasis: Transduction and Integration*. Frontiers in Neuroscience CRC Press/Taylor & Francis.
- De Miguel C, Das S, Lund H, & Mattson DL (2010). T lymphocytes mediate hypertension and kidney damage in Dahl salt-sensitive rats. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* 298: 1136-1142.
- Dennis JM, & Witting PK (2017). Protective role for antioxidants in acute kidney disease. *Nutrients* 9: 1-25.

- Devarajan P (2006). Update on mechanisms of ischemic acute kidney injury. *Journal of the American Society of Nephrology* 17: 1503-1520.
- Doi K, Ishizu T, Tsukamoto-Sumida M, Hiruma T, Yamashita T, Ogasawara E, *et al.* (2014). The high-mobility group protein B1-Toll-like receptor 4 pathway contributes to the acute lung injury induced by bilateral nephrectomy. *Kidney International* 86: 316-326.
- Dong C, Wu Y, Yao J, Wang Y, Yu Y, Rychahou PG, *et al.* (2012). G9a interacts with Snail and is critical for Snail-mediated E-cadherin repression in human breast cancer. *J Clinical Investigation* 122: 1469-1486.
- Dong G, Luo J, Kumar V, & Dong Z (2010). Inhibitors of histone deacetylases suppress cisplatin-induced p53 activation and apoptosis in renal tubular cells. *American Journal of Physiology-Renal Physiology* 298: F293-F300.
- Dong G, Wang L, Wang C-Y, Yang T, Kumar MV, & Dong Z (2008). Induction of apoptosis in renal tubular cells by histone deacetylase inhibitors, a family of anticancer agents. *Journal of Pharmacology and Experimental Therapeutics* 325: 978-984.
- Donnahoo KK, Shames BD, Harken AH, & Meldrum DR (1999). Review article: the role of tumor necrosis factor in renal ischemia-reperfusion injury. *The Journal of Urology* 162: 196-203.
- Donoghue M, Hsieh F, Baronas E, Godbout K, Gosselin M, Stagliano N, *et al.* (2000). A novel angiotensin-converting enzyme-related carboxypeptidase (ACE2) converts angiotensin I to angiotensin 1-9. *Circulation Research* 87: 1-9.
- Donoghue M, Wakimoto H, Maguire CT, Acton S, Hales P, Stagliano N, *et al.* (2003). Heart block, ventricular tachycardia, and sudden death in ACE2 transgenic mice with downregulated connexins. *Journal of Molecular and Cellular Cardiology* 35: 1043-1053.
- Dressler GR (2008). Epigenetics, development, and the kidney. *Journal of the American Society of Nephrology* 19: 2060-2067.

- Dube S, Matam T, Yen J, Mang HE, Dagher PC, Hato T, *et al.* (2017). Endothelial STAT3 modulates protective mechanisms in a mouse ischemia-reperfusion model of acute kidney injury. *Journal of Immunology Research* 2017: 4609502.
- Edelstein CL, Ling H, & Schrier RW (1997). The nature of renal cell injury. *Kidney International* 51: 1341-1351.
- Efrati S, Berman S, Hamad RA, Siman-Tov Y, Ilgiyaev E, Maslyakov I, *et al.* (2011). Effect of captopril treatment on recuperation from ischemia/reperfusion-induced acute renal injury. *Nephrology Dialysis Transplantation* 27: 136-145.
- ElAlfy MS, Ebeid FSE, Kamal TM, Eissa DS, Ismail EAR, & Mohamed SH (2018). Angiotensinogen M235T Gene Polymorphism is a Genetic Determinant of Cerebrovascular and Cardiopulmonary Morbidity in Adolescents with Sickle Cell Disease. *Journal of Stroke and Cerebrovascular Diseases* 28: 441-449.
- Ellefson DD, Espinoza T, Roda N, Maldonado S, & Rodgers KE (2004). Synergistic effects of co-administration of angiotensin 1-7 and Neupogen on hematopoietic recovery in mice. *Cancer Chemotherapy and Pharmacology* 53: 15-24.
- Elmore S (2007). Apoptosis: a review of programmed cell death. *Toxicologic Pathology* 35: 495-516.
- Esteban V, Heringer-Walther S, Sterner-Kock A, de Bruin R, van den Engel S, Wang Y, *et al.* (2009). Angiotensin-(1-7) and the g protein-coupled receptor MAS are key players in renal inflammation. *PLoS One* 4: e5406.
- Etelvino GM, Peluso AAB, & Santos RAS (2014). New Components of the Renin-Angiotensin System: Alamandine and the Mas-Related G Protein-Coupled Receptor D. *Current Hypertension Reports* 16: 1-6.
- Etienne I, Magalhães LVB, Cardoso SA, de Freitas RB, de Oliveira GP, Palotás A, *et al.* (2019). Oxidative stress markers in cognitively intact patients with diabetic neuropathy. *Brain Research Bulletin* 150: 196-200.

- Evans CE, Miners JS, Piva G, Willis CL, Heard DM, Kidd EJ, *et al.* (2020). ACE2 activation protects against cognitive decline and reduces amyloid pathology in the Tg2576 mouse model of Alzheimer's disease. *Acta Neuropathologica* 139: 485-502.
- Evans KJ, & Greenberg A (2005). Hyperkalemia: a review. *Journal of Intensive Care Medicine* 20: 272-290.
- Fadillioglu E, Kurcer Z, Parlakpinar H, Iraz M, & Gursul C (2008). Melatonin treatment against remote organ injury induced by renal ischemia reperfusion injury in diabetes mellitus. *Archives Pharmacal Research* 31: 705-712.
- Fan H, Yang H-C, You L, Wang Y-Y, He W-J, & Hao C-M (2013). The histone deacetylase, SIRT1, contributes to the resistance of young mice to ischemia/reperfusion-induced acute kidney injury. *Kidney International* 83: 404-413.
- Fellows R, Denizot J, Stellato C, Cuomo A, Jain P, Stoyanova E, *et al.* (2018). Microbiota derived short chain fatty acids promote histone crotonylation in the colon through histone deacetylases. *Nature Communications* 9: 1-15.
- Ferenbach DA, & Bonventre JV (2015). Mechanisms of maladaptive repair after AKI leading to accelerated kidney ageing and CKD. *Nature reviews Nephrology* 11: 264-276.
- Fischel T, Hermesh H, Aizenberg D, Zemishlany Z, Munitz H, Benjamini Y, *et al.* (2001). Cyproheptadine versus propranolol for the treatment of acute neuroleptic-induced akathisia: a comparative double-blind study. *Journal of Clinical Psychopharmacology* 21: 612-615.
- Fontecha-Barriuso M, Martin-Sanchez D, Ruiz-Andres O, Poveda J, Sanchez-Niño MD, Valino-Rivas L, *et al.* (2018). Targeting epigenetic DNA and histone modifications to treat kidney disease. *Nephrology Dialysis Transplantation* 33: 1875-1886.
- Fouad AA, & Jresat I (2013). Captopril and telmisartan treatments attenuate cadmium-induced testicular toxicity in rats. *Fundamental and Clinical Pharmacology* 27: 152-160.
- Fraser CL, & Arief AI (1988). Nervous system complications in uremia. *Annals of Internal Medicine* 109: 143-153.

- Fu Y, Tang C, Cai J, Chen G, Zhang D, & Dong Z (2018). Rodent models of AKI-CKD transition. American Journal of Physiology-Renal Physiology 315: F1098-F1106.
- Fuchtemeier M, Brinckmann MP, Foddis M, Kunz A, Po C, Curato C, *et al.* (2015). Vascular change and opposing effects of the angiotensin type 2 receptor in a mouse model of vascular cognitive impairment. Journal of Cerebral Blood Flow & Metabolism 35: 476-484.
- Gadhia S, Shrimp JH, Meier JL, McGee JE, & Dahlin JL (2018). Histone acetyltransferase assays in drug and chemical probe discovery. In Assay Guidance Manual. Eli Lilly & Company and the National Center for Advancing Translational Sciences 1-41.
- Gandolfo MT, Jang HR, Bagnasco SM, Ko GJ, Agreda P, Satpute SR, *et al.* (2009). Foxp3+ regulatory T cells participate in repair of ischemic acute kidney injury. Kidney International 76: 717-729.
- Gao C, Chen G, Liu L, Li X, He J, Jiang L, *et al.* (2013). Impact of high glucose and proteasome inhibitor MG132 on histone H2A and H2B ubiquitination in rat glomerular mesangial cells. Journal of Diabetes Research 2013: 1-10.
- Gao G, Wang W, Tadagavadi RK, Briley NE, Love MI, Miller BA, *et al.* (2014). TRPM2 mediates ischemic kidney injury and oxidant stress through RAC1. The Journal of Clinical Investigation 124: 4989-5001.
- Ghose AK, Viswanadhan VN, & Wendoloski JJ (1999). A knowledge-based approach in designing combinatorial or medicinal chemistry libraries for drug discovery. 1. A qualitative and quantitative characterization of known drug databases. Journal of Combinatorial Chemistry 1: 55-68.
- Golab F, Kadkhodaee M, Zahmatkesh M, Hedayati M, Arab H, Schuster R, *et al.* (2009). Ischemic and non-ischemic acute kidney injury cause hepatic damage. Kidney International 75: 783-792.

- Gong DJ, Wang L, Yang YY, Zhang JJ, & Liu XH (2019). Diabetes aggravates renal ischemia and reperfusion injury in rats by exacerbating oxidative stress, inflammation, and apoptosis. *Renal Failure* 41: 750-761.
- Gonzalez L, Novoa U, Moya J, Gabrielli L, Jalil JE, Garcia L, *et al.* (2018). Angiotensin-(1-9) reduces cardiovascular and renal inflammation in experimental renin-independent hypertension. *Biochemical Pharmacology* 156: 357-370.
- Gorelik G, Carbone LA, & Scicli AG (1998). Angiotensin 1-7 induces bradykinin-mediated relaxation in porcine coronary artery. *The Journal of Pharmacology and Experimental Therapeutics* 286: 403-410.
- Goru SK, Kadakol A, & Gaikwad AB (2017a). Hidden targets of ubiquitin proteasome system: to prevent diabetic nephropathy. *Pharmacological Research* 120: 170-179.
- Goru SK, Kadakol A, Malek V, Pandey A, Sharma N, & Gaikwad AB (2017b). Diminazene acetate prevents nephropathy by increasing glomerular ACE2 and AT2 receptor expression in a rat model of type 1 diabetes. *British Journal of Pharmacology* 174: 3118-3130.
- Goru SK, Kadakol A, Pandey A, Malek V, Sharma N, & Gaikwad AB (2016). Histone H2AK119 and H2BK120 mono-ubiquitination modulate SET7/9 and SUV39H1 in type 1 diabetes-induced renal fibrosis. *Biochemical Journal* 473: 3937-3949.
- Gould SE, Day M, Jones SS, & Dorai H (2002). BMP-7 regulates chemokine, cytokine, and hemodynamic gene expression in proximal tubule cells. *Kidney International* 61: 51-60.
- Grams ME, & Rabb H (2012). The distant organ effects of acute kidney injury. *Kidney International* 81: 942-948.
- Greenway SE, Pack AT, & Greenway FL (1995). Treatment of depression with cyproheptadine. *Pharmacotherapy: The Journal of Human Pharmacology and Drug Therapy* 15: 357-360.

- Greer EL, & Shi Y (2012). Histone methylation: a dynamic mark in health, disease and inheritance. *Nature Reviews Genetics* 13: 343-357.
- Gregoretti I, Lee Y-M, & Goodson HV (2004). Molecular evolution of the histone deacetylase family: functional implications of phylogenetic analysis. *Journal of Molecular Biology* 338: 17-31.
- Greiner D, Bonaldi T, Eskeland R, Roemer E, & Imhof A (2005). Identification of a specific inhibitor of the histone methyltransferase SU (VAR) 3-9. *Nature Chemical Biology* 1: 143-145.
- Griendling KK, Sorescu D, & Ushio-Fukai M (2000). NAD (P) H oxidase: role in cardiovascular biology and disease. *Circulation Research* 86: 494-501.
- Guerra C, Linde-Zwirble WT, & Wunsch H (2012). Risk factors for dementia after critical illness in elderly Medicare beneficiaries. *Critical Care* 16: R233.
- Guo C, Dong G, Liang X, & Dong Z (2019). Epigenetic regulation in AKI and kidney repair: mechanisms and therapeutic implications. *Nature Reviews Nephrology* 15: 220-239.
- Guo Q, Li X, Han H, Li C, Liu S, Gao W, *et al.* (2016). Histone Lysine Methylation in TGF- β 1 Mediated p21 Gene Expression in Rat Mesangial Cells. *Biomed Reserach International* 2016: 6927234.
- Gupta A, Rhodes GJ, Berg DT, Gerlitz B, Molitoris BA, & Grinnell BW (2007). Activated protein C ameliorates LPS-induced acute kidney injury and downregulates renal INOS and angiotensin 2. *American Journal of Physiology-Renal Physiology* 293: 245-254.
- Habibey R, Ajami M, Hesami A, & Pazoki-Toroudi H (2008). The mechanism of preventive effect of captopril on renal ischemia reperfusion injury is independent of ATP dependent potassium channels. *Iran Biomedical Journal* 12: 241-245.
- Hakam AC, Siddiqui AH, & Hussain T (2006). Renal angiotensin II AT2 receptors promote natriuresis in streptozotocin-induced diabetic rats. *American Journal of Physiology-Renal Physiology* 290: F503-F508.

- Hammad F, Al-Salam S, & Lubbad L (2013). 609 Renoprotective effects of aliskiren following ischemia reperfusion injury. European Urology Supplements 12: e609.
- Han SJ, Kim J-H, Kim JI, & Park KM (2016). Inhibition of microtubule dynamics impedes repair of kidney ischemia/reperfusion injury and increases fibrosis. Scientific Reports 6: 27775.
- Han SJ, & Lee HT (2019). Mechanisms and therapeutic targets of ischemic acute kidney injury. Kidney Research and Clinical Practice. 38: 427-440.
- Han W, Waikar S, Johnson A, Betensky R, Dent C, Devarajan P, *et al.* (2008). Urinary biomarkers in the early diagnosis of acute kidney injury. Kidney International 73: 863-869.
- Hang LH, Xu ZK, Wei SY, Shu WW, Luo H, & Chen J (2017). Spinal SET 7/9 may contribute to the maintenance of cancer-induced bone pain in mice. Clinical and Experimental Pharmacology and Physiology 44: 1001-1007.
- Hansen MK, Gammelager H, Jacobsen C-J, Hjortdal VE, Layton JB, Rasmussen BS, *et al.* (2015). Acute kidney injury and long-term risk of cardiovascular events after cardiac surgery: a population-based cohort study. Journal of Cardiothoracic and Vascular Anesthesia 29: 617-625.
- Harel Z, Gilbert C, Wald R, Bell C, Perl J, Juurlink D, *et al.* (2012). The effect of combination treatment with aliskiren and blockers of the renin-angiotensin system on hyperkalaemia and acute kidney injury: systematic review and meta-analysis. Bmj 344: e42.
- Hasegawa K, Wakino S, Yoshioka K, Tatematsu S, Hara Y, Minakuchi H, *et al.* (2010). Kidney-specific overexpression of Sirt1 protects against acute kidney injury by retaining peroxisome function. Journal of Biological Chemistry 285: 13045-13056.
- Havasi A, Haegele JA, Gall JM, Blackmon S, Ichimura T, Bonegio RG, *et al.* (2013). Histone acetyl transferase (HAT) HBO1 and JADE1 in epithelial cell regeneration. American Journal of Pathology 182: 152-162.

- He S, Liu N, Bayliss G, & Zhuang S (2013). EGFR activity is required for renal tubular cell dedifferentiation and proliferation in a murine model of folic acid-induced acute kidney injury. *American Journal of Physiology-Renal Physiology* 304: F356-366.
- He X, Zhang T, Tolosa M, Goru SK, Chen X, Misra PS, *et al.* (2019). A new, easily generated mouse model of diabetic kidney fibrosis. *Scientific Reports* 9: 1-12.
- Herget-Rosenthal S, Marggraf G, Hüsing J, Goring F, Pietruck F, Janssen O, *et al.* (2004). Early detection of acute renal failure by serum cystatin C. *Kidney International* 66: 1115-1122.
- Heringer-Walther S, Eckert K, Schumacher S-M, Uharek L, Wulf-Goldenberg A, Gembardt F, *et al.* (2009). Angiotensin-(1-7) stimulates hematopoietic progenitor cells in vitro and in vivo. *Haematologica*: 94(6):857-60.
- Hershko A, Ciechanover A, & Varshavsky A (2000). Basic Medical Research Award. The ubiquitin system. *Nature Medicine* 6: 1073-1081.
- Hertzberg D, Sartipy U, & Holzmann MJ (2015). Type 1 and type 2 diabetes mellitus and risk of acute kidney injury after coronary artery bypass grafting. *American Heart Journal* 170: 895-902.
- Hewitson TD, Holt SG, Tan S-J, Wigg B, Samuel CS, & Smith ER (2017). Epigenetic modifications to H3K9 in renal tubulointerstitial cells after unilateral ureteric obstruction and TGF- β 1 stimulation. *Frontiers in Pharmacology* 8: 307.
- Higuchi S, Ohtsu H, Suzuki H, Shirai H, Frank GD, & Eguchi S (2007). Angiotensin II signal transduction through the AT1 receptor: novel insights into mechanisms and pathophysiology. *Clinical Science* 112: 417-428.
- Hirano T, Fujiwara T, Niwa H, Hirano M, Ohira K, Okazaki Y, *et al.* (2018). Development of Novel Inhibitors for Histone Methyltransferase SET7/9 based on Cyproheptadine. *ChemMedChem* 13:1530-1540.
- Holderied A, & Anders HJ (2014). Animal models of kidney inflammation in translational medicine. *Drug Discovery Today: Disease Models* 11: 19-27.

- Holderied A, Kraft F, Marschner JA, Weidenbusch M, & Anders H-J (2020). “Point of no return” in unilateral renal ischemia reperfusion injury in mice. *Journal of Biomedical Science* 27: 1-15.
- Hollands C, Bartolotti N, & Lazarov O (2016). Alzheimer's disease and hippocampal adult neurogenesis; exploring shared mechanisms. *Frontiers in Neuroscience* 10: 178.
- Holliday R (1987). The inheritance of epigenetic defects. *Science* 238: 163-170.
- Hoste EA, Kellum JA, Selby NM, Zarbock A, Palevsky PM, Bagshaw SM, *et al.* (2018). Global epidemiology and outcomes of acute kidney injury. *Nature Reviews Nephrology* 4: 607-625.
- Hrenak J, Arendasova K, Rajkovicova R, Aziriova S, Repova K, Krajcirovicova K, *et al.* (2013). Protective effect of captopril, olmesartan, melatonin and compound 21 on doxorubicin-induced nephrotoxicity in rats. *Physiological Research* 62: S181.
- Hrenak J, Paulis L, & Simko F (2016). Angiotensin a/alamandine/mrgd axis: another clue to understanding cardiovascular pathophysiology. *International Journal of Molecular Sciences* 17: 1-9.
- Huang J, Dorsey J, Chuikov S, Zhang X, Jenuwein T, Reinberg D, *et al.* (2010). G9a and Glp methylate lysine 373 in the tumor suppressor p53. *Journal of Biological Chemistry* 285: 9636-9641.
- Huang J, Wan D, Li J, Chen H, Huang K, & Zheng L (2015a). Histone acetyltransferase PCAF regulates inflammatory molecules in the development of renal injury. *Epigenetics* 10: 62-71.
- Huang K-P, Chen C, Hao J, Huang J-Y, Liu P-Q, & Huang H-Q (2015b). AGEs-RAGE system down-regulates Sirt1 through the ubiquitin-proteasome pathway to promote FN and TGF- β 1 expression in male rat glomerular mesangial cells. *Endocrinology* 156: 268-279.
- Hursh BE, Ronsley R, Islam N, Mammen C, & Panagiotopoulos C (2017). Acute kidney injury in children with type 1 diabetes hospitalized for diabetic ketoacidosis. *JAMA pediatrics* 171: e170020-e170020.

- Husain-Syed F, Rosner MH, & Ronco C (2019). Distant organ dysfunction in acute kidney injury. *Acta Physiologica*: 1-10.
- Ichiki T, Labosky PA, Shiota C, Okuyama S, Imagawa Y, Fogo A, *et al.* (1995). Effects on blood pressure and exploratory behaviour of mice lacking angiotensin II type-2 receptor. *Nature* 377: 748-750.
- Irifuku T, Doi S, Sasaki K, Doi T, Nakashima A, Ueno T, *et al.* (2016). Inhibition of H3K9 histone methyltransferase G9a attenuates renal fibrosis and retains klotho expression. *Kidney International* 89: 147-157.
- Isaka Y (2016). Epidermal growth factor as a prognostic biomarker in chronic kidney diseases. *Annals of Translational Medicine* 4: 1-4.
- Isobe-Sasaki Y, Fukuda M, Ogiyama Y, Sato R, Miura T, Fuwa D, *et al.* (2017). Sodium balance, circadian BP rhythm, heart rate variability, and intrarenal renin–angiotensin–aldosterone and dopaminergic systems in acute phase of ARB therapy. *Physiological Reports* 5: 1-13.
- Jackman HL, Massad MG, Sekosan M, Tan F, Brovkovich V, Marcic BM, *et al.* (2002). Angiotensin 1-9 and 1-7 release in human heart: role of cathepsin A. *Hypertension* 39: 976-981.
- Jang H-S, Han SJ, Kim JI, Lee S, Lipschutz JH, & Park KM (2013). Activation of ERK accelerates repair of renal tubular epithelial cells, whereas it inhibits progression of fibrosis following ischemia/reperfusion injury. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease* 1832: 1998-2008.
- Jang HR, Gandolfo MT, Ko GJ, Satpute SR, Racusen L, & Rabb H (2010). B cells limit repair after ischemic acute kidney injury. *Journal of the American Society of Nephrology* 21: 654-665.
- Jang HR, & Rabb H (2009). The innate immune response in ischemic acute kidney injury. *Clinical Immunology* 130: 41-50.

- Jha JC, Banal C, Chow BS, Cooper ME, & Jandeleit-Dahm K (2016). Diabetes and kidney disease: role of oxidative stress. *Antioxid Redox Signal* 25: 657-684.
- Jorres A, Gahl GM, Dobis C, Polenakovic MH, Cakalaroski K, Rutkowski B, *et al.* (1999). Haemodialysis-membrane biocompatibility and mortality of patients with dialysis-dependent acute renal failure: a prospective randomised multicentre trial. *The Lancet* 354: 1337-1341.
- Joseph JP, Mecca AP, Regenhardt RW, Bennion DM, Rodríguez V, Desland F, *et al.* (2014). The angiotensin type 2 receptor agonist Compound 21 elicits cerebroprotection in endothelin-1 induced ischemic stroke. *Neuropharmacology* 81: 134-141.
- Kadakol A, Malek V, Goru SK, Pandey A, & Gaikwad AB (2015). Esculetin reverses histone H2A/H2B ubiquitination, H3 dimethylation, acetylation and phosphorylation in preventing type 2 diabetic cardiomyopathy. *Journal of Functional Foods* 17: 127-136.
- Kako K, Kato M, Matsuoka T, & Mustapha A (1988). Depression of membrane-bound Na⁺-K⁺-ATPase activity induced by free radicals and by ischemia of kidney. *American Journal of Physiology-Cell Physiology* 254: C330-C337.
- Kakoki M, McGarrah RW, Kim H-S, & Smithies O (2007). Bradykinin B1 and B2 receptors both have protective roles in renal ischemia/reperfusion injury. *Proceedings of the National Academy of Sciences* 104: 7576-7581.
- Kamel AS, Abdelkader NF, Abd El-Rahman SS, Emara M, Zaki HF, & Khattab MM (2018). Stimulation of ACE2/ANG(1-7)/Mas Axis by Diminazene Ameliorates Alzheimer's Disease in the D-Galactose-Ovariectomized Rat Model: Role of PI3K/Akt Pathway. *Molecular Neurobiology* 55: 8188-8202.
- Kanagasundaram NS (2015). Pathophysiology of ischaemic acute kidney injury. *Annals of Clinical Biochemistry* 52: 193-205.
- Kanasaki M, Nagai T, Kitada M, Koya D, & Kanasaki K (2011). Elevation of the antifibrotic peptide N-acetyl-seryl-aspartyl-lysyl-proline: a blood pressure-independent

beneficial effect of angiotensin I-converting enzyme inhibitors. *Fibrogenesis & Tissue Repair* 4: 25.

Kangussu LM, Almeida-Santos AF, Moreira FA, Fontes MA, Santos RA, Aguiar DC, *et al.* (2017). Reduced anxiety-like behavior in transgenic rats with chronically overproduction of angiotensin-(1–7): Role of the Mas receptor. *Behavioural Brain Research* 331: 193-198.

Karnik SS, Unal H, Kemp JR, Tirupula KC, Eguchi S, Vanderheyden PM, *et al.* (2015). International Union of Basic and Clinical Pharmacology. XCIX. Angiotensin Receptors: Interpreters of Pathophysiological Angiotensinergic Stimuli. *Pharmacological Reviews* 67: 754-819.

Karpe PA, Gupta J, Marthong RF, Ramarao P, & Tikoo K (2012). Insulin resistance induces a segmental difference in thoracic and abdominal aorta: differential expression of AT1 and AT2 receptors. *Journal of Hypertension* 30: 132-146.

Karpe PA, & Tikoo K (2014). Heat shock prevents insulin resistance-induced vascular complications by augmenting angiotensin-(1-7) signaling. *Diabetes* 63: 1124-1139.

Kaschina E, Grzesiak A, Li J, Foryst-Ludwig A, Timm M, Rompe F, *et al.* (2008). Angiotensin II type 2 receptor stimulation: a novel option of therapeutic interference with the renin-angiotensin system in myocardial infarction? *Circulation* 118: 2523-2532.

Kashani K, Al-Khafaji A, Ardiles T, Artigas A, Bagshaw SM, Bell M, *et al.* (2013). Discovery and validation of cell cycle arrest biomarkers in human acute kidney injury. *Critical Care* 17: 1-12.

Kaushal GP, & Shah SV (2014). Challenges and advances in the treatment of AKI. *Journal of the American Society of Nephrology* 25: 877-883.

Kellum JA (2016). Why are patients still getting and dying from acute kidney injury? *Current Opinion in Critical Care* 22: 513-519.

Kelly K (2003). Distant effects of experimental renal ischemia/reperfusion injury. *Journal of the American Society of Nephrology* 14: 1549-1558.

- Kezic A, Stajic N, & Thaiss F (2017). Innate immune response in kidney ischemia/reperfusion injury: potential target for therapy. *Journal of Immunology Research* 2017.
- Khastar H (2015). Protective effects of vitamin E against liver damage caused by renal ischemia reperfusion. *Renal Failure* 37: 494-496.
- Kim J, & Padanilam BJ (2015). Renal denervation prevents long-term sequelae of ischemic renal injury. *Kidney International* 87: 350-358.
- Kim JI, Jung KJ, Jang H-S, & Park KM (2013). Gender-specific role of HDAC11 in kidney ischemia-and reperfusion-induced PAI-1 expression and injury. *American Journal of Physiology-Renal Physiology* 305: F61-F70.
- Kim M, Park SW, Kim M, D'Agati VD, & Lee HT (2010). Isoflurane activates intestinal sphingosine kinase to protect against bilateral nephrectomy-induced liver and intestine dysfunction. *American Journal of Physiology-Renal Physiology* 300: F167-F176.
- Kim N, Jung Y, Nam M, Kang MS, Lee MK, Cho Y, *et al.* (2017). Angiotensin II affects inflammation mechanisms via AMPK-related signalling pathways in HL-1 atrial myocytes. *Scientific Repts* 7: 1-12.
- Kim SM, Kim YG, Jeong KH, Lee SH, Lee TW, Ihm CG, *et al.* (2012). Angiotensin II-induced mitochondrial Nox4 is a major endogenous source of oxidative stress in kidney tubular cells. *PloS One* 7: 60-70.
- Kinsey GR, & Okusa MD (2012). Role of leukocytes in the pathogenesis of acute kidney injury. *Critical Care* 16: 1-5.
- Kobori H, Harrison-Bernard LM, & Navar LG (2002). Urinary excretion of angiotensinogen reflects intrarenal angiotensinogen production. *Kidney International* 61: 579-585.

- Kobori H, Nangaku M, Navar LG, & Nishiyama A (2007). The intrarenal renin-angiotensin system: from physiology to the pathobiology of hypertension and kidney disease. *Pharmacological Reviews* 59: 251-287.
- Kocak C, Kocak FE, Akcilar R, Bayat Z, Aras B, Metineren MH, *et al.* (2016). Effects of captopril, telmisartan and bardoxolone methyl (CDDO-Me) in ischemia-reperfusion-induced acute kidney injury in rats: an experimental comparative study. *Clinical and Experimental Pharmacology and Physiology* 43: 230-241.
- Kokkonen JO, Saarinen J, & Kovanen PT (1997). Regulation of local angiotensin II formation in the human heart in the presence of interstitial fluid. Inhibition of chymase by protease inhibitors of interstitial fluid and of angiotensin-converting enzyme by Ang-(1-9) formed by heart carboxypeptidase A-like activity. *Circulation* 95: 1455-1463.
- Kontogiannis J, & Burns KD (1998). Role of AT1 angiotensin II receptors in renal ischemic injury. *American Journal of Physiology-Renal Physiology* 274: 79-90.
- Kooistra SM, & Helin K (2012). Molecular mechanisms and potential functions of histone demethylases. *Nature Reviews Molecular Cell Biology* 13: 297-311.
- Kornberg RD (1974). Chromatin structure: a repeating unit of histones and DNA. *Science* 184: 868-871.
- Koulis C, Chow BS, McKelvey M, Steckelings UM, Unger T, Thallas-Bonke V, *et al.* (2015). AT2R agonist, compound 21, is reno-protective against type 1 diabetic nephropathy. *Hypertension* 65: 1073-1081.
- Kouzarides T (2007). Chromatin modifications and their function. *Cell* 128: 693-705.
- Kovalčíková A, Gyurászová M, Vavrinčová-Yaghi D, Vavrinec P, Tóthová L, Boor P, *et al.* (2018). Oxidative stress in the brain caused by acute kidney injury. *Metabolic Brain Disease* 33: 961-967.
- Kramer AA, Postler G, Salhab KF, Mendez C, Carey LC, & Rabb H (1999). Renal ischemia/reperfusion leads to macrophage-mediated increase in pulmonary vascular permeability. *Kidney International* 55: 2362-2367.

- Kramkowski K, Mogielnicki A, Leszczynska A, & Buczko W (2010). Angiotensin-(1-9), the product of angiotensin I conversion in platelets, enhances arterial thrombosis in rats. *Journal of Physiology and Pharmacology : an Official Journal of the Polish Physiological Society*, 61: 317-324.
- Krasaelap A, & Madani S (2017). Cyproheptadine: a potentially effective treatment for functional gastrointestinal disorders in children. *Pediatric Annals* 46: e120-e125.
- Kuhad A, Pilkhwal S, Sharma S, Tirkey N, & Chopra K (2007). Effect of curcumin on inflammation and oxidative stress in cisplatin-induced experimental nephrotoxicity. *Journal of Agricultural and Food Chemistry* 55: 10150-10155.
- Kurash JK, Lei H, Shen Q, Marston WL, Granda BW, Fan H, *et al.* (2008). Methylation of p53 by Set7/9 mediates p53 acetylation and activity in vivo. *Molecular Cell* 29: 392-400.
- Kuriakose S, & Uzonna JE (2014). Diminazene aceturate (Berenil), a new use for an old compound? *International Immunopharmacology* 21: 342-345.
- Kwon T-H, Nielsen J, Kim YH, Knepper MA, Frokiær J, & Nielsen S (2003). Regulation of sodium transporters in the thick ascending limb of rat kidney: response to angiotensin II. *American Journal of Physiology-Renal Physiology* 285: 152-165.
- Lai Y, Chen J, Tsai H, Chen T, & Hung W (2015). The SUV39H1 inhibitor chaetocin induces differentiation and shows synergistic cytotoxicity with other epigenetic drugs in acute myeloid leukemia cells. *Blood Cancer Journal* 5: e313-e313.
- Lai Y, Deng J, Wang M, Wang M, Zhou L, Meng G, *et al.* (2019). Vagus nerve stimulation protects against acute liver injury induced by renal ischemia reperfusion via antioxidant stress and anti-inflammation. *Biomedicine & Pharmacotherapy* 117: 109062.
- Lane K, Dixon JJ, MacPhee IA, & Philips BJ (2013). Renohepatic crosstalk: does acute kidney injury cause liver dysfunction? *Nephrology Dialysis Transplantation* 28: 1634-1647.

- Lautner RQ, Villela DC, Fraga-Silva RA, Silva N, Verano-Braga T, Costa-Fraga F, *et al.* (2013). Discovery and Characterization of AlamandineNovelty and Significance: A Novel Component of the Renin-Angiotensin System. *Circulation Research* 112: 1104-1111.
- Lazzeri E, Angelotti ML, Conte C, Anders H-J, & Romagnani P (2019). Surviving acute organ failure: cell polyplloidization and progenitor proliferation. *Trends in Molecular medicine*, 25: 366-381.
- Lazzeri E, Angelotti ML, Peired A, Conte C, Marschner JA, Maggi L, *et al.* (2018). Endocycle-related tubular cell hypertrophy and progenitor proliferation recover renal function after acute kidney injury. *Nature Communications* 9: 1-18.
- Le Clef N, Verhulst A, D'Haese PC, & Vervaet BA (2016). Unilateral renal ischemia-reperfusion as a robust model for acute to chronic kidney injury in mice. *PloS one* 11: e0152153.
- Le Dorze M, Legrand M, Payen D, & Ince C (2009). The role of the microcirculation in acute kidney injury. *Current Opinion in Critical Care* 15: 503-508.
- Lea-Henry TN, Carland JE, Stocker SL, Sevastos J, & Roberts DM (2018). Clinical Pharmacokinetics in Kidney Disease: Fundamental Principles. *Clinical Journal of the American Society of Nephrology* 13: 1085-1095.
- Leao N, Etelvino G, & Santos R (2016). Abstract P281: Alamandine-induced Vasorelaxation is Selectively Increased in SP-SHR Hypertension. 2016 68: AP281.
- Lech M, Avila-Ferrufino A, Allam R, Segerer S, Khandoga A, Krombach F, *et al.* (2009). Resident dendritic cells prevent postischemic acute renal failure by help of single Ig IL-1 receptor-related protein. *The Journal of Immunology* 183: 4109-4118.
- Lee D, Faubel S, & Edelstein C (2011). Cytokines in acute kidney injury (AKI). *Clinical Nephrology* 76: 165-173.
- Lee JC, Park JH, Kim IH, Cho GS, Ahn JH, Tae HJ, *et al.* (2017a). Neuroprotection of ischemic preconditioning is mediated by thioredoxin 2 in the hippocampal CA1 region following a subsequent transient cerebral ischemia. *Brain Pathology* 27: 276-291.

- Lee SA, Cozzi M, Bush EL, & Rabb H (2018). Distant organ dysfunction in acute kidney injury: a review. *American Journal of Kidney Diseases* 72: 846-856.
- Lee SA, Noel S, Sadasivam M, Hamad AR, & Rabb H (2017b). Role of immune cells in acute kidney injury and repair. *Nephron* 137: 282-286.
- Lekawannijit S, Kompa AR, & Krum H (2016). Protein-bound uremic toxins: a long overlooked culprit in cardiorenal syndrome. *American Journal of Physiology-Renal Physiology* 311: F52-F62.
- Leonard MO, Kieran NE, Howell K, Burne MJ, Varadarajan R, Dhakshinamoorthy S, *et al.* (2006). Reoxygenation-specific activation of the antioxidant transcription factor Nrf2 mediates cytoprotective gene expression in ischemia-reperfusion injury. *FASEB Journal* 20: 2624-2626.
- Levine M, Wang Z, Bhatti T, Wang Y, Aufhauser D, McNeal S, *et al.* (2015). Class-specific histone/protein deacetylase inhibition protects against renal ischemia reperfusion injury and fibrosis formation. *American Journal of Transplantation* 15: 965-973.
- Li L, Huang L, Sung S-SJ, Vergis AL, Rosin DL, Rose Jr CE, *et al.* (2008a). The chemokine receptors CCR2 and CX3CR1 mediate monocyte/macrophage trafficking in kidney ischemia-reperfusion injury. *Kidney International* 74: 1526-1537.
- Li L, McBride DW, Doycheva D, Dixon BJ, Krafft PR, Zhang JH, *et al.* (2015). G-CSF attenuates neuroinflammation and stabilizes the blood-brain barrier via the PI3K/Akt/GSK-3 β signaling pathway following neonatal hypoxia-ischemia in rats. *Experimental Neurology* 272: 135-144.
- Li L, Zhang Y, Xu M, Rong R, Wang J, & Zhu T (2016a). Inhibition of histone methyltransferase EZH2 ameliorates early acute renal allograft rejection in rats. *BMC Immunology* 17: 41.
- Li LX, Fan LX, Zhou JX, Grantham JJ, Calvet JP, Sage J, *et al.* (2017). Lysine methyltransferase SMYD2 promotes cyst growth in autosomal dominant polycystic kidney disease. *The Journal of Clinical Investigation* 127: 2751-2764.

- Li X, Li C, Li X, Cui P, Li Q, Guo Q, *et al.* (2016b). Involvement of Histone Lysine Methylation in p21 Gene Expression in Rat Kidney In Vivo and Rat Mesangial Cells In Vitro under Diabetic Conditions. *Journal of Diabetes Research* 2016: 3853242.
- Li Y, Reddy MA, Miao F, Shanmugam N, Yee J-K, Hawkins D, *et al.* (2008b). Role of the histone H3 lysine 4 methyltransferase, SET7/9, in the regulation of NF-κB-dependent inflammatory genes relevance to diabetes and inflammation. *Journal of Biological Chemistry* 283: 26771-26781.
- Liang H, Huang Q, Liao M-j, Xu F, Zhang T, He J, *et al.* (2019). EZH2 plays a crucial role in ischemia/reperfusion-induced acute kidney injury by regulating p38 signaling. *Inflammation Research* 68: 325-336.
- Liano F, Junco E, Madero R, Pascual J, & Verde E (1998). The spectrum of acute renal failure in the intensive care unit compared with that seen in other settings. *Kidney International-Supplements*: 16-24.
- Lin SH, Ho WT, Wang YT, Chuang CT, Chuang LY, & Guh JY (2016). Histone methyltransferase Suv39h1 attenuates high glucose-induced fibronectin and p21WAF1 in mesangial cells. *The International Journal of Biochemistry & Cell Biology* 78: 96-105.
- Liu F, Barsyte-Lovejoy D, Li F, Xiong Y, Korboukh V, Huang X-P, *et al.* (2013a). Discovery of an in vivo chemical probe of the lysine methyltransferases G9a and GLP. *Journal of Medicinal Chemistry* 56: 8931-8942.
- Liu L, Mao L, Wu X, Wu T, Liu W, Yang Y, *et al.* (2019a). BRG1 regulates endothelial-derived IL-33 to promote ischemia-reperfusion induced renal injury and fibrosis in mice. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease* 1865: 2551-2561.
- Liu L, Zou J, Guan Y, Zhang Y, Zhang W, Zhou X, *et al.* (2019b). Blocking the histone lysine 79 methyltransferase DOT1L alleviates renal fibrosis through inhibition of renal fibroblast activation and epithelial-mesenchymal transition. *The FASEB Journal* 33: 11941-11958.

- Liu M, Grigoryev DN, Crow MT, Haas M, Yamamoto M, Reddy SP, *et al.* (2009). Transcription factor Nrf2 is protective during ischemic and nephrotoxic acute kidney injury in mice. *Kidney International* 76: 277-285.
- Liu M, Liang Y, Chigurupati S, Lathia JD, Pletnikov M, Sun Z, *et al.* (2008). Acute kidney injury leads to inflammation and functional changes in the brain. *Journal of the American Society of Nephrology* 19: 1360-1370.
- Liu N, He S, Ma L, Ponnusamy M, Tang J, Tolbert E, *et al.* (2013b). Blocking the class I histone deacetylase ameliorates renal fibrosis and inhibits renal fibroblast activation via modulating TGF-beta and EGFR signaling. *PloS one* 8: e54001-e54001.
- Lizakowski S, Tylicki L, Renke M, Rutkowski P, Heleniak Z, Ślawińska-Morawska M, *et al.* (2012). Effect of aliskiren on proteinuria in non-diabetic chronic kidney disease: a double-blind, crossover, randomised, controlled trial. *International Urology and Nephrology* 44: 1763-1770.
- Locksley RM, Killeen N, & Lenardo MJ (2001). The TNF and TNF receptor superfamilies: integrating mammalian biology. *Cell* 104: 487-501.
- Lopez B, Salom MG, Arregui B, Valero F, & Fenoy FJ (2003). Role of superoxide in modulating the renal effects of angiotensin II. *Hypertension* 42: 1150-1156.
- Lu R, Kiernan MC, Murray A, Rosner MH, & Ronco C (2015). Kidney–brain crosstalk in the acute and chronic setting. *Nature Reviews Nephrology* 11: 707.
- Luo Q-h, Chen M-l, Sun F-j, Chen Z-l, Li M-y, Zeng W, *et al.* (2014). KIM-1 and NGAL as biomarkers of nephrotoxicity induced by gentamicin in rats. *Molecular and Cellular Biochemistry* 397: 53-60.
- Lv Q, Yang Q, Cui Y, Yang J, Wu G, Liu M, *et al.* (2017). Effects of Taurine on ACE, ACE2 and HSP70 Expression of Hypothalamic-Pituitary-Adrenal Axis in Stress-Induced Hypertensive Rats. *Advances in Experimental Medicine and Biology* 975 Pt 2: 871-886.

- Ma Z, Wei Q, Dong G, Huo Y, & Dong Z (2014). DNA damage response in renal ischemia–reperfusion and ATP-depletion injury of renal tubular cells. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease* 1842: 1088-1096.
- Macconi D, Sangalli F, Bonomelli M, Conti S, Condorelli L, Gagliardini E, *et al.* (2009). Podocyte repopulation contributes to regression of glomerular injury induced by ACE inhibition. *The American Journal of Pathology* 174: 797-807.
- Macedo LM, da Silva Souza ÁP, De Maria MLdA, Borges CL, de Almeida Soares CM, Pedrino GR, *et al.* (2016). Cardioprotective effects of diminazene aceturate in pressure-overloaded rat hearts. *Life Sciences* 155: 63-69.
- Machado-Silva A, Passos-Silva D, Santos RA, & Sinisterra RD (2016). Therapeutic uses for Angiotensin-(1-7). *Expert Opinion on Therapeutic Patents* 26: 669-678.
- Macia Heras M, Del Castillo Rodriguez N, & Navarro Gonzalez J (2012). The renin–angiotensin–aldosterone system in renal and cardiovascular disease and the effects of its pharmacological blockade. *Journal of Diabetes and Metabolism* 3: 1-24.
- Mackie FE, Campbell DJ, & Meyer TW (2001). Intrarenal angiotensin and bradykinin peptide levels in the remnant kidney model of renal insufficiency. *Kidney International* 59: 1458-1465.
- Majumder S, Thieme K, Batchu SN, Alghamdi TA, Bowskill BB, Kabir MG, *et al.* (2018). Shifts in podocyte histone H3K27me3 regulate mouse and human glomerular disease. *The Journal of Clinical Investigation* 128: 483-499.
- Makris K, & Spanou L (2016). Acute kidney injury: definition, pathophysiology and clinical phenotypes. *The Clinical Biochemist Reviews* 37: 85-98.
- Malek M, & Nematbakhsh M (2014). The preventive effects of diminazene aceturate in renal ischemia/reperfusion injury in male and female rats. *Advances in Preventive Medicine* 2014: 740647.
- Malek M, & Nematbakhsh M (2015). Renal ischemia/reperfusion injury; from pathophysiology to treatment. *Journal of Renal Injury Prevention* 4: 20-27.

- Malek V, & Gaikwad AB (2017). Neprilysin inhibitors: a new hope to halt the diabetic cardiovascular and renal complications? *Biomedicine & Pharmacotherapy* 90: 752-759.
- Malek V, & Gaikwad AB (2019). Telmisartan and thiophan combination treatment attenuates fibrosis and apoptosis in preventing diabetic cardiomyopathy. *Cardiovascular Research* 115: 373-384.
- Malek V, Sharma N, & Gaikwad AB (2019a). Simultaneous inhibition of neprilysin and activation of ACE2 prevented diabetic cardiomyopathy. *Pharmacological Reports* 71: 958-967.
- Malek V, Sharma N, Sankrityayan H, & Gaikwad AB (2019b). Concurrent neprilysin inhibition and renin-angiotensin system modulations prevented diabetic nephropathy. *Life Sciences* 221: 159-167.
- Mar D, Gharib SA, Zager RA, Johnson A, Denisenko O, & Bomsztyk K (2015). Heterogeneity of epigenetic changes at ischemia/reperfusion-and endotoxin-induced acute kidney injury genes. *Kidney International* 88: 734-744.
- Marenzi G, Cosentino N, Milazzo V, De Metrio M, Rubino M, Campodonico J, *et al.* (2018). Acute kidney injury in diabetic patients with acute myocardial infarction: Role of acute and chronic glycemia. *Journal of the American Heart Association* 7: e008122.
- Marko L, Vigolo E, Hinze C, Park J-K, Roel G, Balogh A, *et al.* (2016). Tubular epithelial NF-κB activity regulates ischemic AKI. *Journal of the American Society of Nephrology* 27: 2658-2669.
- Marre M, Bernadet P, Gallois Y, Savagner F, Guyene TT, Hallab M, *et al.* (1994). Relationships between angiotensin I converting enzyme gene polymorphism, plasma levels, and diabetic retinal and renal complications. *Diabetes* 43: 384-388.
- Marre M, Jeunemaitre X, Gallois Y, Rodier M, Chatellier G, Sert C, *et al.* (1997). Contribution of genetic polymorphism in the renin-angiotensin system to the development of renal complications in insulin-dependent diabetes: Genetique de la Nephropathie

- Diabetique (GENEDIAB) study group. The Journal of Clinical Investigation 99: 1585-1595.
- Martin JL, Gruszczak AV, Beach TE, Murphy MP, & Saeb-Parsy K (2019). Mitochondrial mechanisms and therapeutics in ischaemia reperfusion injury. Pediatric Nephrology 34: 1167-1174.
- Martinez A-M, & Cavalli G (2006). The role of polycomb group proteins in cell cycle regulation during development. Cell Cycle 5: 1189-1197.
- Marumo T, Hishikawa K, Yoshikawa M, & Fujita T (2008). Epigenetic regulation of BMP7 in the regenerative response to ischemia. Journal of the American Society of Nephrology 19: 1311-1320.
- Matavelli LC, Huang J, & Siragy HM (2011). Angiotensin AT(2) receptor stimulation inhibits early renal inflammation in renovascular hypertension. Hypertension 57: 308-313.
- Matejovic M, Ince C, Chawla LS, Blantz R, Molitoris BA, Rosner MH, *et al.* (2016). Renal hemodynamics in AKI: in search of new treatment targets. Journal of American Society of Nephrology 27: 49-58.
- McCarthy CA, Vinh A, Miller AA, Hallberg A, Alterman M, Callaway JK, *et al.* (2014). Direct angiotensin AT2 receptor stimulation using a novel AT2 receptor agonist, compound 21, evokes neuroprotection in conscious hypertensive rats. PloS One 9: e95762.
- McCord JM (1985). Oxygen-derived free radicals in postischemic tissue injury. New England Journal of Medicine 312: 159-163.
- McCullough JW, Renner B, & Thurman JM The role of the complement system in acute kidney injury. Seminars in Nephrology 33: 543-556.
- McKinney CA, Fattah C, Loughrey CM, Milligan G, & Nicklin SA (2014). Angiotensin-(1-7) and angiotensin-(1-9): function in cardiac and vascular remodelling. Clinical Science 126: 815-827.

- McKinney CA, Kennedy S, Baillie G, Milligan G, & Nicklin SA (2015). 178 Angiotensin-(1–9) inhibits vascular smooth muscle cell proliferation and migration in vitro and neointimal formation in vivo BMJ Publishing Group Ltd and British Cardiovascular Society Heart 2015 101: A101.
- McLaughlin N, Wang F, Saifudeen Z, & El-Dahr SS (2014). In situ histone landscape of nephrogenesis. *Epigenetics* 9: 222-235.
- McMahon BA, & Koyner JL (2016). Risk stratification for acute kidney injury: Are biomarkers enough? *Advances in Chronic Kidney Disease* 23: 167-178.
- Meersch M, Schmidt C, Hoffmeier A, Van Aken H, Wempe C, Gerss J, *et al.* (2017). Prevention of cardiac surgery-associated AKI by implementing the KDIGO guidelines in high risk patients identified by biomarkers: the PrevAKI randomized controlled trial. *Intensive Care Medicine* 43: 1551-1561.
- Mehrotra P, Patel JB, Ivancic CM, Collett JA, & Basile DP (2015). Th-17 cell activation in response to high salt following acute kidney injury is associated with progressive fibrosis and attenuated by AT-1R antagonism. *Kidney International* 88: 776-784.
- Mendez CE, Der Mesropian PJ, Mathew RO, & Slawski B (2016). Hyperglycemia and acute kidney injury during the perioperative period. *Current Diabetes Reports* 16: 10.
- Metzger R, Bader M, Ludwig T, Berberich C, Bunnemann B, & Ganter D (1995). Expression of the mouse and rat mas proto-oncogene in the brain and peripheral tissues. *FEBS letters* 357: 27-32.
- Mimura I, Tanaka T, & Nangaku M (2016). New insights into molecular mechanisms of epigenetic regulation in kidney disease. *Clinical and Experimental Pharmacology and Physiology* 43: 1159-1167.
- Min J, Feng Q, Li Z, Zhang Y, & Xu R-M (2003). Structure of the catalytic domain of human DOT1L, a non-SET domain nucleosomal histone methyltransferase. *Cell* 112: 711-723.

- Min LJ, Mogi M, Tsukuda K, Jing F, Ohshima K, Nakaoka H, *et al.* (2014). Direct stimulation of angiotensin II type 2 receptor initiated after stroke ameliorates ischemic brain damage. *American Journal of Hypertension* 27: 1036-1044.
- Mishra J, Ma Q, Prada A, Mitsnefes M, Zahedi K, Yang J, *et al.* (2003). Identification of neutrophil gelatinase-associated lipocalin as a novel early urinary biomarker for ischemic renal injury. *Journal of American Society of Nephrology* 14: 2534-2543.
- Miura S-i, Matsuo Y, Kiya Y, Karnik SS, & Saku K (2010). Molecular mechanisms of the antagonistic action between AT1 and AT2 receptors. *Biochemical and Biophysical Research Communications* 391: 85-90.
- Mohammadi M, Najafi H, Yarjani ZM, Vaezi G, & Hojati V (2019). Piperine pretreatment attenuates renal ischemia-reperfusion induced liver injury. *Heliyon* 5: e02180.
- Molinas SM, Cortes-Gonzalez C, Gonzalez-Bobadilla Y, Monasterolo LA, Cruz C, Elías MM, *et al.* (2009). Effects of losartan pretreatment in an experimental model of ischemic acute kidney injury. *Nephron Experimental Nephrology* 112: e10-e19.
- Moreno JA, Izquierdo MC, Sanchez-Niño MD, Suárez-Alvarez B, Lopez-Larrea C, Jakubowski A, *et al.* (2011). The inflammatory cytokines TWEAK and TNF α reduce renal klotho expression through NF κ B. *Journal of the American Society of Nephrology* 22: 1315-1325.
- Mori J, Patel VB, Ramprasath T, Alrob OA, DesAulniers J, Scholey JW, *et al.* (2014). Angiotensin 1-7 mediates renoprotection against diabetic nephropathy by reducing oxidative stress, inflammation, and lipotoxicity. *American Journal of Physiology-Renal Physiology* 306: F812-F821.
- Moriyama T, Tsuruta Y, Kojima C, Itabashi M, Sugiura H, Takei T, *et al.* (2012). Beneficial effect of aliskiren combined with olmesartan in reducing urinary protein excretion in patients with chronic kidney disease. *International Urology and Nephrology* 44: 841-845.

- Munoz MC, Burghi V, Miquet JG, Cervino IA, Quiroga DT, Mazziotta L, *et al.* (2017). Chronic blockade of the AT2 receptor with PD123319 impairs insulin signaling in C57BL/6 mice. *Peptides* 88: 37-45.
- Munshi R, Johnson A, Siew ED, Ikitler TA, Ware LB, Wurfel MM, *et al.* (2011). MCP-1 gene activation marks acute kidney injury. *Journal of the American Society of Nephrology* 22: 165-175.
- Murphy T, Alexander RW, Griendling KK, Runge MS, & Bernstein KE (1991). Isolation of a cDNA encoding the vascular type-1 angiotensin II receptor. *Nature* 351: 233-236.
- Murray KC, Nakae A, Stephens MJ, Rank M, D'amico J, Harvey PJ, *et al.* (2010). Recovery of motoneuron and locomotor function after spinal cord injury depends on constitutive activity in 5-HT 2C receptors. *Nature medicine* 16: 694.
- Naito M, Bomsztyk K, & Zager RA (2008). Endotoxin mediates recruitment of RNA polymerase II to target genes in acute renal failure. *Journal of the American Society of Nephrology* 19: 1321-1330.
- Naito M, Bomsztyk K, & Zager RA (2009a). Renal ischemia-induced cholesterol loading: transcription factor recruitment and chromatin remodeling along the HMG CoA reductase gene. *The American Journal of Pathology* 174: 54-62.
- Naito M, Zager RA, & Bomsztyk K (2009b). BRG1 increases transcription of proinflammatory genes in renal ischemia. *Journal of the American Society of Nephrology* 20: 1787-1796.
- Nakazawa D, Kumar SV, Marschner J, Desai J, Holderied A, Rath L, *et al.* (2017). Histones and neutrophil extracellular traps enhance tubular necrosis and remote organ injury in ischemic AKI. *Journal of the American Society of Nephrology* 28: 1753-1768.
- Namsolleck P, Recarti C, Foulquier S, Steckelings UM, & Unger T (2014). AT(2) receptor and tissue injury: therapeutic implications. *Current Hypertension Reports* 16: 1-10.

- Ni J, Hou X, Wang X, Shi Y, Xu L, Zheng X, *et al.* (2019). 3-deazaneplanocin A protects against cisplatin-induced renal tubular cell apoptosis and acute kidney injury by restoration of E-cadherin expression. *Cell Death & Disease* 10: 1-15.
- Nieuwenhuijs-Moeke GJ, Pischke SE, Berger SP, Sanders JSF, Pol RA, Struys MM, *et al.* (2020). Ischemia and reperfusion injury in kidney transplantation: relevant mechanisms in injury and repair. *Journal of Clinical Medicine* 9: 253.
- Nongnuch A, Panorchan K, & Davenport A (2014). Brain–kidney crosstalk. *Critical care* 18: 1-11.
- Novitskaya T, McDermott L, Zhang KX, Chiba T, Pauksakon P, Hukriede NA, *et al.* (2014). A PTBA small molecule enhances recovery and reduces postinjury fibrosis after aristolochic acid-induced kidney injury. *American Journal of Physiology-Renal Physiology* 306: F496-F504.
- Ocaranza MP, Godoy I, Jalil JE, Varas M, Collantes P, Pinto M, *et al.* (2006). Enalapril attenuates downregulation of angiotensin-converting enzyme 2 in the late phase of ventricular dysfunction in myocardial infarcted rat. *Hypertension* 48: 572-578.
- Ocaranza MP, & Jalil JE (2012). Protective role of the ACE2/Ang-(1-9) axis in cardiovascular remodeling. *International Journal of Hypertension* 2012: 1-12.
- Ocaranza MP, Moya J, Barrientos V, Alzamora R, Hevia D, Morales C, *et al.* (2014). Angiotensin-(1-9) reverses experimental hypertension and cardiovascular damage by inhibition of the angiotensin converting enzyme/Ang II axis. *Journal of Hypertension* 32: 771-783.
- Oezkur M, Wagner M, Weismann D, Krannich JH, Schimmer C, Riegler C, *et al.* (2015). Chronic hyperglycemia is associated with acute kidney injury in patients undergoing CABG surgery—a cohort study. *BMC Cardiovascular Disorders* 15: 1-6.
- Ohshima K, Mogi M, Jing F, Iwanami J, Tsukuda K, Min LJ, *et al.* (2012). Direct angiotensin II type 2 receptor stimulation ameliorates insulin resistance in type 2 diabetes mice with PPARgamma activation. *PloS one* 7: e48387.

- Oktaba K, Gutiérrez L, Gagneur J, Girardot C, Sengupta AK, Furlong EE, *et al.* (2008). Dynamic regulation by polycomb group protein complexes controls pattern formation and the cell cycle in Drosophila. *Developmental Cell* 15: 877-889.
- Okuma H, Iijima K, Yasuda T, Tokuoka K, & Kitagawa Y (2013). Preventive effect of cyproheptadine hydrochloride in refractory patients with frequent migraine. *Springerplus* 2: 1-4.
- Omoniwa D, Ogunpolu B, Omoniwa AM, Adedokun R, Olaifa A, Oyagbemi A, *et al.* (2019). Effect of Diminazene aceturate treatment on some cardiovascular parameters and antioxidant status of apparently healthy horses. *The FASEB Journal* 33: 819.811-819.811.
- Ono M, Sakao Y, Tsuji T, Ohashi N, Yasuda H, Nishiyama A, *et al.* (2015). Role of intrarenal (pro) renin receptor in ischemic acute kidney injury in rats. *Clinical and Experimental Nephrology* 19: 185-196.
- Ortiz A, Lorz C, Catalán MP, Danoff TM, Yamasaki Y, Egido J, *et al.* (2000). Expression of apoptosis regulatory proteins in tubular epithelium stressed in culture or following acute renal failure. *Kidney International* 57: 969-981.
- Oudit GY, Liu GC, Zhong J, Basu R, Chow FL, Zhou J, *et al.* (2010). Human recombinant ACE2 reduces the progression of diabetic nephropathy. *Diabetes* 59: 529-538.
- Pabla N, Huang S, Mi Q-S, Daniel R, & Dong Z (2008). ATR-Chk2 signaling in p53 activation and DNA damage response during cisplatin-induced apoptosis. *Journal of Biological Chemistry* 283: 6572-6583.
- Padda RS, Shi Y, Lo C-S, Zhang S-L, & Chan JS (2015). Angiotensin-(1-7): a novel peptide to treat hypertension and nephropathy in diabetes? *Journal of Diabetes & Metabolism* 6:1-12.
- Palkovits M, Sebekova K, Gallatz K, Boor P, Sebekova Jr K, Klassen A, *et al.* (2009). Neuronal activation in the CNS during different forms of acute renal failure in rats. *Neuroscience* 159: 862-882.

- Paller MS, Manivel JC, Patten M, & Barry M (1992). Prostaglandins protect kidneys against ischemic and toxic injury by a cellular effect. *Kidney International* 42: 1345-1354.
- Pandey A, & Gaikwad AB (2017a). AT2 receptor agonist Compound 21: A silver lining for diabetic nephropathy. *European Journal of Pharmacology* 815: 251-257.
- Pandey A, & Gaikwad AB (2017b). Compound 21 and Telmisartan combination mitigates type 2 diabetic nephropathy through amelioration of caspase mediated apoptosis. *Biochemical and Biophysical Research Communications* 487: 827-833.
- Pandey A, Goru SK, Kadakol A, Malek V, & Gaikwad AB (2015). Differential regulation of angiotensin converting enzyme 2 and nuclear factor- κ B by angiotensin II receptor subtypes in type 2 diabetic kidney. *Biochimie* 118: 71-81.
- Pandey A, Raj P, Goru SK, Kadakol A, Malek V, Sharma N, *et al.* (2017). Esculetin ameliorates hepatic fibrosis in high fat diet induced non-alcoholic fatty liver disease by regulation of FoxO1 mediated pathway. *Pharmacological Reports* 69: 666-672.
- Pang M, Kothapally J, Mao H, Tolbert E, Ponnusamy M, Chin YE, *et al.* (2009). Inhibition of histone deacetylase activity attenuates renal fibroblast activation and interstitial fibrosis in obstructive nephropathy. *American Journal of Physiology-Renal Physiology* 297: F996-F1005.
- Panico K, Abrahao MV, Trentin-Sonoda M, Muzzi-Filho H, Vieyra A, & Carneiro-Ramos MS (2019). Cardiac Inflammation after Ischemia-Reperfusion of the Kidney: Role of the Sympathetic Nervous System and the Renin-Angiotensin System. *Cellular Physiology and Biochemistry* 53: 587-605.
- Pardridge WM (2005). The blood-brain barrier: Bottleneck in brain drug development. *NeuroRX* 2: 3-14.
- Parikh CR, Jani A, Melnikov VY, Faubel S, & Edelstein CL (2004). Urinary interleukin-18 is a marker of human acute tubular necrosis1. *American Journal of Kidney Disease* 43: 405-414.

- Park JK, Mervaala EM, Muller DN, Menne J, Fiebeler A, Luft FC, *et al.* (2009). Rosuvastatin protects against angiotensin II-induced renal injury in a dose-dependent fashion. *Journal of Hypertension* 27: 599-605.
- Park SW, Chen SW, Kim M, Brown KM, Kolls JK, D D'Agati V, *et al.* (2011). Cytokines induce small intestine and liver injury after renal ischemia or nephrectomy. *Laboratory Investigation* 91: 63-84.
- Park SW, Kim M, Kim JY, Ham A, Brown KM, Mori-Akiyama Y, *et al.* (2012). Paneth cell-mediated multiorgan dysfunction after acute kidney injury. *The Journal of Immunology* 189: 5421-5433.
- Patschan D, & Müller G (2016). Acute kidney injury in diabetes mellitus. *International Journal of Nephrology* 2016: 1-7.
- Patschan D, Schwarze K, Lange A, Meise N, Henze E, Becker JU, *et al.* (2013). Bone morphogenetic protein-5 and early endothelial outgrowth cells (eEOCs) in acute ischemic kidney injury (AKI) and 5/6-chronic kidney disease. *American Journal of Physiology-Renal Physiology* 305: F314-F322.
- Pazoki-Toroudi HR, Hesami A, Vahidi S, Sahebjam F, Seifi B, & Djahanguiri B (2003). The preventive effect of captopril or enalapril on reperfusion injury of the kidney of rats is independent of angiotensin II AT1 receptors. *Fundamental & Clinical Pharmacology* 17: 595-598.
- Peng J, Li X, Zhang D, Chen J-K, Su Y, Smith SB, *et al.* (2015). Hyperglycemia, p53, and mitochondrial pathway of apoptosis are involved in the susceptibility of diabetic models to ischemic acute kidney injury. *Kidney International* 87: 137-150.
- Peppicelli S, Bianchini F, Contena C, Tombaccini D, & Calorini L (2013). Acidic pH via NF-κB favours VEGF-C expression in human melanoma cells. *Clinical & Experimental Metastasis* 30: 957-967.
- Persson PB (2003). Renin: origin, secretion and synthesis. *The Journal of Physiology* 552: 667-671.

- Philips BJ, Lane K, Dixon J, & MacPhee I (2014). The effects of acute renal failure on drug metabolism. *Expert Opinion on Drug Metabolism & Toxicology* 10: 11-23.
- Piantadosi CA, & Suliman HB (2006). Mitochondrial transcription factor A induction by redox activation of nuclear respiratory factor 1. *Journal of Biological Chemistry* 281: 324-333.
- Pickart CM (2001). Mechanisms underlying ubiquitination. *Annual Review of Biochemistry* 70: 503-533.
- Pickering JW, & Endre ZH (2014). The definition and detection of acute kidney injury. *Journal of Renal Injury Prevention* 3: 21-25.
- Pires-Luís AS, Vieira-Coimbra M, Vieira FQ, Costa-Pinheiro P, Silva-Santos R, Dias PC, *et al.* (2015). Expression of histone methyltransferases as novel biomarkers for renal cell tumor diagnosis and prognostication. *Epigenetics* 10: 1033-1043.
- Popovic D, Vucic D, & Dikic I (2014). Ubiquitination in disease pathogenesis and treatment. *Nature Medicine* 20: 1242-1253.
- Porrello ER, D'amore A, Curl CL, Allen AM, Harrap SB, Thomas WG, *et al.* (2009). Angiotensin II type 2 receptor antagonizes angiotensin II type 1 receptor-mediated cardiomyocyte autophagy. *Hypertension* 53: 1032-1040.
- Portela A, & Esteller M (2010). Epigenetic modifications and human disease. *Nature Biotechnology* 28: 1057.
- Portilla D, Dent C, Sugaya T, Nagothu K, Kundi I, Moore P, *et al.* (2008). Liver fatty acid-binding protein as a biomarker of acute kidney injury after cardiac surgery. *Kidney International* 73: 465-472.
- Quiroga DT, Muñoz MC, Gil C, Pfleifer M, Toblli JE, Steckelings UM, *et al.* (2018). Chronic administration of the angiotensin type 2 receptor agonist C21 improves insulin sensitivity in C57 BL/6 mice. *Physiological Reports* 6: e13824.

- Rabb H, Griffin MD, McKay DB, Swaminathan S, Pickkers P, Rosner MH, *et al.* (2016). Inflammation in AKI: Current Understanding, Key Questions, and Knowledge Gaps. *Journal of American Society of Nephrology* 27: 371-379.
- Rafiq K, Hitomi H, Nakano D, & Nishiyama A (2011). Pathophysiological roles of aldosterone and mineralocorticoid receptor in the kidney. *Journal of Pharmacological Sciences* 115: 1-7.
- Rai NK, Tripathi K, Sharma D, & Shukla VK (2005). Apoptosis: a basic physiologic process in wound healing. *The International Journal of Lower Extremity Wounds* 4: 138-144.
- Rajapaksha IG, Mak KY, Huang P, Burrell LM, Angus PW, & Herath CB (2018). The small molecule drug diminazene aceturate inhibits liver injury and biliary fibrosis in mice. *Scientific Reports* 8: 10175.
- Ranganathan P, Hamad R, Mohamed R, Jayakumar C, Muthusamy T, & Ramesh G (2016). Histone deacetylase-mediated silencing of AMWAP expression contributes to cisplatin nephrotoxicity. *Kidney International* 89: 317-326.
- Ratliff BB, Abdulmahdi W, Pawar R, & Wolin MS (2016). Oxidant mechanisms in renal injury and disease. *Antioxidants & Redox Signaling* 25: 119-146.
- Raup-Konsavage WM, Wang Y, Wang WW, Feliers D, Ruan H, & Reeves WB (2018). Neutrophil peptidyl arginine deiminase-4 has a pivotal role in ischemia/reperfusion-induced acute kidney injury. *Kidney International* 93: 365-374.
- Reddy MA, & Natarajan R (2015). Recent developments in epigenetics of acute and chronic kidney diseases. *Kidney International* 88: 250-261.
- Reddy MA, Sumanth P, Lanting L, Yuan H, Wang M, Mar D, *et al.* (2014). Losartan reverses permissive epigenetic changes in renal glomeruli of diabetic db/db mice. *Kidney International* 85: 362-373.

- Reed JC, Zha H, Aime-Sempe C, Takayama S, & Wang H-G (1996). Structure—Function Analysis of Bcl-2 Family Proteins. In Mechanisms of lymphocyte activation and immune regulation VI. Springer, pp 99-112.
- Reule S, & Gupta S (2011). Kidney regeneration and resident stem cells. *Organogenesis* 7: 135-139.
- Rigat B, Hubert C, Alhenc-Gelas F, Cambien F, Corvol P, & Soubrier F (1990). An insertion/deletion polymorphism in the angiotensin I-converting enzyme gene accounting for half the variance of serum enzyme levels. *The Journal of Clinical Investigation* 86: 1343-1346.
- Rodriguez Romo R, Benitez K, Barrera Chimal J, Perez Villalva R, Gomez A, Aguilar Leon D, *et al.* (2016). AT1 receptor antagonism before ischemia prevents the transition of acute kidney injury to chronic kidney disease. *Kidney International* 89: 363-373.
- Rompe F, Artuc M, Hallberg A, Alterman M, Stroder K, Thone-Reineke C, *et al.* (2010). Direct angiotensin II type 2 receptor stimulation acts anti-inflammatory through epoxyeicosatrienoic acid and inhibition of nuclear factor kappaB. *Hypertension* 55: 924-931.
- Ronco C (2008). Cardiorenal and renocardiac syndromes: clinical disorders in search of a systematic definition. *The International Journal of Artificial Organs* 31: 1-2.
- Ronco C, Haapio M, House AA, Anavekar N, & Bellomo R (2008). Cardiorenal syndrome. *Journal of the American College of Cardiology* 52: 1527-1539.
- Ross MJ, & Nangaku M (2017). ACE2 as therapy for glomerular disease: the devil is in the detail. *Kidney International* 91: 1269-1271.
- Rossetto D, Avvakumov N, & Côté J (2012). Histone phosphorylation: a chromatin modification involved in diverse nuclear events. *Epigenetics* 7: 1098-1108.
- Ruiz-Andres O, Sanchez-Nino MD, Cannata-Ortiz P, Ruiz-Ortega M, Egido J, Ortiz A, *et al.* (2016a). Histone lysine crotonylation during acute kidney injury in mice. *Disease Models & Mechanisms* 9: 633-645.

- Ruiz-Andres O, Suarez-Alvarez B, Sánchez-Ramos C, Monsalve M, Sanchez-Niño MD, Ruiz-Ortega M, *et al.* (2016b). The inflammatory cytokine TWEAK decreases PGC-1 α expression and mitochondrial function in acute kidney injury. *Kidney International* 89: 399-410.
- Ruiz-Ortega M, Esteban V, Suzuki Y, Ruperez M, Mezzano S, Ardiles L, *et al.* (2003). Renal expression of angiotensin type 2 (AT2) receptors during kidney damage. *Kidney International* 64: S21-S26.
- Ruthenburg AJ, Allis CD, & Wysocka J (2007). Methylation of lysine 4 on histone H3: intricacy of writing and reading a single epigenetic mark. *Molecular Cell* 25: 15-30.
- Sakao Y, Kato A, Tsuji T, Yasuda H, Togawa A, Fujigaki Y, *et al.* (2011). Cisplatin induces Sirt1 in association with histone deacetylation and increased Werner syndrome protein in the kidney. *Clinical and Experimental Nephrology* 15: 363-372.
- Salahudeen AK, Haider N, & May W (2004). Cold ischemia and the reduced long-term survival of cadaveric renal allografts. *Kidney International* 65: 713-718.
- Salama M, Mohamed Farrag S, Mohamed Amin M, Abdel-Aziz Ali A, Sheashaa H, Sobh M, *et al.* (2013). Up-regulation of TLR-4 in the brain after ischemic kidney-induced encephalopathy in the rat. *CNS & Neurological Disorders-Drug Targets (Formerly Current Drug Targets-CNS & Neurological Disorders)* 12: 583-586.
- Sanchez-Pozos K, Barrera-Chimal J, Garzon-Muvdi J, Perez-Villalva R, Rodriguez-Romo R, Cruz C, *et al.* (2012). Recovery from ischemic acute kidney injury by spironolactone administration. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association* 27: 3160-3169.
- Santos RA, Castro CH, Gava E, Pinheiro SV, Almeida AP, de Paula RD, *et al.* (2006). Impairment of in vitro and in vivo heart function in angiotensin-(1-7) receptor MAS knockout mice. *Hypertension* 47: 996-1002.
- Santos RA, Ferreira AJ, & Simoes ESAC (2008). Recent advances in the angiotensin-converting enzyme 2-angiotensin(1-7)-Mas axis. *Experimental Physiology* 93: 519-527.

- Santos RA, Simoes e Silva AC, Maric C, Silva DM, Machado RP, de Buhr I, *et al.* (2003). Angiotensin-(1-7) is an endogenous ligand for the G protein-coupled receptor Mas. *Proceedings of the National Academy of Sciences of the United States of America* 100: 8258-8263.
- Santos RAS, Sampaio WO, Alzamora AC, Motta-Santos D, Alenina N, Bader M, *et al.* (2018). The ACE2/Angiotensin-(1-7)/MAS Axis of the Renin-Angiotensin System: Focus on Angiotensin-(1-7). *Physiological Reviews* 98: 505-553.
- Santos SH, Giani JF, Burghi V, Miquet JG, Qadri F, Braga JF, *et al.* (2014). Oral administration of angiotensin-(1-7) ameliorates type 2 diabetes in rats. *Journal of molecular medicine (Berlin, Germany)* 92: 255-265.
- Sasaki K, Doi S, Nakashima A, Irifuku T, Yamada K, Kokoroishi K, *et al.* (2016). Inhibition of SET domain-containing lysine methyltransferase 7/9 ameliorates renal fibrosis. *Journal of the American Society of Nephrology* 27: 203-215.
- Sasaki K, Yamano Y, Bardhan S, Iwai N, Murray JJ, Hasegawa M, *et al.* (1991). Cloning and expression of a complementary DNA encoding a bovine adrenal angiotensin II type-1 receptor. *Nature* 351: 230-233.
- Savergnini SQ, Beiman M, Lautner RQ, de Paula-Carvalho V, Allahdadi K, Pessoa DC, *et al.* (2010). Vascular relaxation, antihypertensive effect, and cardioprotection of a novel peptide agonist of the MAS receptor. *Hypertension* 56: 112-120.
- Sawhney S, & Fraser SD (2017). Epidemiology of AKI: utilizing large databases to determine the burden of AKI. *Advances in Chronic Kidney Disease* 24: 194-204.
- Sayed-Tabatabaei F, Oostra B, Isaacs A, Van Duijn C, & Witteman J (2006). ACE polymorphisms. *Circulation Research* 98: 1123-1133.
- Scalia R, Gong Y, Berzins B, Freund B, Feather D, Landesberg G, *et al.* (2011). A Novel Role for Calpain in the Endothelial Dysfunction Induced by Activation of Angiotensin II

- Type 1 Receptor Signaling Novelty and Significance. *Circulation Research* 108: 1102-1111.
- Scholpa N, Zhang X, Kolli R, & Cummings B (2014). Epigenetic changes in p21 expression in renal cells after exposure to bromate. *Toxicological Sciences* 141: 432-440.
- Schoolwerth AC, Sica DA, Ballermann BJ, & Wilcox CS (2001). Renal considerations in angiotensin converting enzyme inhibitor therapy: a statement for healthcare professionals from the Council on the Kidney in Cardiovascular Disease and the Council for High Blood Pressure Research of the American Heart Association. *Circulation* 104: 1985-1991.
- Schweizer S, Harms C, Lerch H, Flynn J, Hecht J, Yildirim F, *et al.* (2015). Inhibition of histone methyltransferases SUV39H1 and G9a leads to neuroprotection in an in vitro model of cerebral ischemia. *Journal of Cerebral Blood Flow & Metabolism* 35: 1640-1647.
- Sehnine M, Ferhat M, Sena S, Gombert J, Goujon J, Thierry A, *et al.* (2018). IL-33 receptor ST2 deficiency attenuates renal ischaemia-reperfusion injury in euglycaemic, but not streptozotocin-induced hyperglycaemic mice. *Diabetes & Metabolism* 44: 55-60.
- Seifi B, Kadkhodaee M, Najafi A, & Mahmoudi A (2014). Protection of liver as a remote organ after renal ischemia-reperfusion injury by renal ischemic postconditioning. *International Journal of Nephrology* 2014: 1-4.
- Serteser M, Koken T, Kahraman A, Yilmaz K, Akbulut G, & Dilek ON (2002). Changes in hepatic TNF- α levels, antioxidant status, and oxidation products after renal ischemia/reperfusion injury in mice. *Journal of Surgical Research* 107: 234-240.
- Shanmugam N, Kim YS, Lanting L, & Natarajan R (2003). Regulation of cyclooxygenase-2 expression in monocytes by ligation of the receptor for advanced glycation end products. *Journal of Biological Chemistry* 278: 34834-34844.
- Shao X, Tian L, Xu W, Zhang Z, Wang C, Qi C, *et al.* (2014). Diagnostic value of urinary kidney injury molecule 1 for acute kidney injury: a meta-analysis. *PloS One* 9: 389-391.
- Sharfuddin AA, & Molitoris BA (2011). Pathophysiology of ischemic acute kidney injury. *Nature Reviews Nephrology* 7: 189-200.

- Sharma S, & Taliyan R (2016). Epigenetic modifications by inhibiting histone deacetylases reverse memory impairment in insulin resistance induced cognitive deficit in mice. *Neuropharmacology* 105: 285-297.
- Sharma S, Taliyan R, & Ramagiri S (2015). Histone deacetylase inhibitor, trichostatin A, improves learning and memory in high-fat diet-induced cognitive deficits in mice. *Journal of Molecular Neuroscience* 56: 1-11.
- Shen W-C, Chou Y-H, Huang H-P, Sheen J-F, Hung S-C, & Chen H-F (2018). Induced pluripotent stem cell-derived endothelial progenitor cells attenuate ischemic acute kidney injury and cardiac dysfunction. *Stem Cell Research & Therapy* 9: 1-12.
- Shen X, Mizuguchi G, Hamiche A, & Wu C (2000). A chromatin remodelling complex involved in transcription and DNA processing. *Nature* 406: 541-544.
- Shi Y, Xu L, Tang J, Fang L, Ma S, Ma X, *et al.* (2017). Inhibition of HDAC6 protects against rhabdomyolysis-induced acute kidney injury. *American Journal of Physiology-Renal Physiology* 312: F502-F515.
- Shiao CC, Wu PC, Huang TM, Lai TS, Yang WS, Wu CH, *et al.* (2015). Long-term remote organ consequences following acute kidney injury. *Critical Care* 19: 438.
- Shuttleworth VG, Gaughan L, Nawafa L, Mooney CA, Cobb SL, Sheerin NS, *et al.* (2018). The methyltransferase SET9 regulates TGFB1 activation of renal fibroblasts via interaction with SMAD3. *Journal of Cell Science* 131: jcs207761.
- Siddiqi FS, Majumder S, Thai K, Abdalla M, Hu P, Advani SL, *et al.* (2016). The histone methyltransferase enzyme enhancer of zeste homolog 2 protects against podocyte oxidative stress and renal injury in diabetes. *Journal of the American Society of Nephrology* 27: 2021-2034.
- Siew ED, Fissell WH, Tripp CM, Blume JD, Wilson MD, Clark AJ, *et al.* (2017). Acute kidney injury as a risk factor for delirium and coma during critical illness. *American Journal of Respiratory and Critical Care Medicine* 195: 1597-1607.

- Silva AR, Aguilar EC, Alvarez-Leite JI, da Silva RF, Arantes RM, Bader M, *et al.* (2013). Mas receptor deficiency is associated with worsening of lipid profile and severe hepatic steatosis in ApoE-knockout mice. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* 305: R1323-R1330.
- Silver SA, Harel Z, McArthur E, Nash DM, Acedillo R, Kitchlu A, *et al.* (2018). Causes of Death after a Hospitalization with AKI. *Journal of American Society of Nephrology* 29: 1001-1010.
- Sims III RJ, Nishioka K, & Reinberg D (2003). Histone lysine methylation: a signature for chromatin function. *TRENDS in Genetics* 19: 629-639.
- Singh K, Xiao L, Remondino A, Sawyer DB, & Colucci WS (2001). Adrenergic regulation of cardiac myocyte apoptosis. *Journal of Cellular Physiology* 189: 257-265.
- Siragy HM (2007). Angiotensin AT1 and AT2 receptors—the battle for health and disease. *Nephrology Dialysis Transplantation* 22: 3128-3130.
- Skrypnyk NI, Harris RC, & de Caestecker MP (2013). Ischemia-reperfusion model of acute kidney injury and post injury fibrosis in mice. *JoVE (Journal of Visualized Experiments)*: e50495.
- Soler MJ, Riera M, Crespo M, Mir M, Marquez E, Pascual MJ, *et al.* (2012). Circulating angiotensin-converting enzyme 2 activity in kidney transplantation: a longitudinal pilot study. *Nephron Clinical Practice* 121: c144-150.
- Solomon SD, Hee Shin S, Shah A, Skali H, Desai A, Kober L, *et al.* (2011). Effect of the direct renin inhibitor aliskiren on left ventricular remodelling following myocardial infarction with systolic dysfunction. *European Heart Journal* 32: 1227-1234.
- Song N, Thaiss F, & Guo L (2019). NF κ B and Kidney Injury. *Frontiers in Immunology* 10: 815.
- Souza LKM, Nicolau LA, Sousa NA, Araújo TS, Sousa FBM, Costa DS, *et al.* (2016). Diminazene aceturate, an angiotensin-converting enzyme II activator, prevents gastric

- mucosal damage in mice: Role of the angiotensin-(1–7)/Mas receptor axis. *Biochemical Pharmacology* 112: 50-59.
- Srisawat N, Sileanu FE, Murugan R, Bellomo R, Calzavacca P, Cartin-Ceba R, *et al.* (2015). Variation in risk and mortality of acute kidney injury in critically ill patients: a multicenter study. *American Journal of Nephrology* 41: 81-88.
- Steckelings UM, Paulis L, Namsolleck P, & Unger T (2012). AT2 receptor agonists: hypertension and beyond. *Current Opinion in Nephrology and Hypertension* 21: 142-146.
- Stein J, Osgood R, Barnes J, Reineck H, Pinckard R, & Mcmanus LM (1985). The role of complement in the pathogenesis of postischemic acute renal failure. *Mineral and Electrolyte Metabolism* 11: 256-261.
- Su H, Wan C, Song A, Qiu Y, Xiong W, & Zhang C (2019). Oxidative Stress and Renal Fibrosis: Mechanisms and Therapies. In *Renal Fibrosis: Mechanisms and Therapies*. Springer, pp 585-604.
- Sugiyama S, Hanaki Y, Ogawa T, Hieda N, Taki K, & Ozawa T (1988). The effects of SUN 1165, a novel sodium channel blocker, on ischemia-induced mitochondrial dysfunction and leakage of lysosomal enzymes in canine hearts. *Biochemical and Biophysical Research Communications* 157: 433-439.
- Sun D-Q, Zhang L, Zheng C-F, Liu W-Y, Zheng KI, Chen X-M, *et al.* (2019). Metabolic Acidosis in Critically Ill Cirrhotic Patients with Acute Kidney Injury. *Journal of Clinical and Translational Hepatology* 7: 112.
- Sun G, Reddy MA, Yuan H, Lanting L, Kato M, & Natarajan R (2010). Epigenetic histone methylation modulates fibrotic gene expression. *Journal of the American Society of Nephrology* 21: 2069-2080.
- Sun H, Zou S, Candiotti KA, Peng Y, Zhang Q, Xiao W, *et al.* (2017). Octreotide attenuates acute kidney injury after hepatic ischemia and reperfusion by enhancing autophagy. *Scientific Reports* 7: 42701.

- Sun J, Shannon M, Ando Y, Schnackenberg LK, Khan NA, Portilla D, *et al.* (2012). Serum metabolomic profiles from patients with acute kidney injury: a pilot study. *Journal of Chromatography B* 893: 107-113.
- Sung FL, Zhu TY, Au-Yeung KK, Siow YL, & O K (2002). Enhanced MCP-1 expression during ischemia/reperfusion injury is mediated by oxidative stress and NF-kappaB. *Kidney International* 62: 1160-1170.
- Susantitaphong P, Cruz DN, Cerdá J, Abulfaraj M, Alqahtani F, Koulouridis I, *et al.* (2013). World incidence of AKI: a meta-analysis. *Clinical Journal of the American Society of Nephrology* 8: 1482-1493.
- Swaminathan M, Shaw AD, Phillips-Bute BG, McGugan-Clark PL, Archer LE, Talbert S, *et al.* (2007). Trends in acute renal failure associated with coronary artery bypass graft surgery in the United States. *Critical Care Medicine* 35: 2286-2291.
- Tachibana M, Sugimoto K, Fukushima T, & Shinkai Y (2001). Set domain-containing protein, G9a, is a novel lysine-preferring mammalian histone methyltransferase with hyperactivity and specific selectivity to lysines 9 and 27 of histone H3. *Journal of Biological Chemistry* 276: 25309-25317.
- Tachibana M, Sugimoto K, Nozaki M, Ueda J, Ohta T, Ohki M, *et al.* (2002). G9a histone methyltransferase plays a dominant role in euchromatic histone H3 lysine 9 methylation and is essential for early embryogenesis. *Genes & Development* 16: 1779-1791.
- Takahashi N, & Smithies O (2004). Human genetics, animal models and computer simulations for studying hypertension. *Trends in Genetics* 20: 136-145.
- Takemoto Y, Ito A, Niwa H, Okamura M, Fujiwara T, Hirano T, *et al.* (2016). Identification of cyproheptadine as an inhibitor of SET domain containing lysine methyltransferase 7/9 (Set7/9) that regulates estrogen-dependent transcription. *Journal of Medicinal Chemistry* 59: 3650-3660.
- Takiyama Y, & Haneda M (2014). Hypoxia in diabetic kidneys. *BioMed Research International* 2014: 1-10.

- Tamura R, Doi S, Nakashima A, Sasaki K, Maeda K, Ueno T, *et al.* (2018). Inhibition of the H3K4 methyltransferase SET7/9 ameliorates peritoneal fibrosis. *PLoS One* 13: e0196844.
- Tan M, Luo H, Lee S, Jin F, Yang JS, Montellier E, *et al.* (2011). Identification of 67 histone marks and histone lysine crotonylation as a new type of histone modification. *Cell* 146: 1016-1028.
- Tanaka S, & Okusa MD (2020). Crosstalk between the nervous system and the kidney. *Kidney International* 97: 466-476.
- Tanaka S, Tanaka T, Kawakami T, Takano H, Sugahara M, Saito H, *et al.* (2017). Vascular adhesion protein-1 enhances neutrophil infiltration by generation of hydrogen peroxide in renal ischemia/reperfusion injury. *Kidney International* 92: 154-164.
- Tang J, Yan Y, Zhao TC, Gong R, Bayliss G, Yan H, *et al.* (2014). Class I HDAC activity is required for renal protection and regeneration after acute kidney injury. *American Journal of Physiology-Renal Physiology* 307: F303-F316.
- Tang J, & Zhuang S (2015). Epigenetics in acute kidney injury. *Current Opinion in Nephrology and Hypertension* 24: 351-358.
- Tang J, & Zhuang S (2019). Histone acetylation and DNA methylation in ischemia/reperfusion injury. *Clinical Science* 133: 597-609.
- Taskin E, & Guven C (2017). Local Renin-Angiotensin System at Liver and Crosstalk with Hepatic Diseases. In *Renin-Angiotensin System-Past, Present and Future*. IntechOpen, London.
- Tetzner A, Gebolys K, Meinert C, Klein S, Uhlich A, Trebicka J, *et al.* (2016). G-Protein-Coupled Receptor MrgD Is a Receptor for Angiotensin-(1–7) Involving Adenylyl Cyclase, cAMP, and Phosphokinase A Novelty and Significance. *Hypertension* 68: 185-194.

- Thakar CV, Christianson A, Himmelfarb J, & Leonard AC (2011). Acute kidney injury episodes and chronic kidney disease risk in diabetes mellitus. *Clinical Journal of American Society of Nephrology*: p CJN. 01120211.
- Thurman JM, Lucia MS, Ljubanovic D, & Holers VM (2005). Acute tubular necrosis is characterized by activation of the alternative pathway of complement. *Kidney International* 67: 524-530.
- Tikoo K, Lau SS, & Monks TJ (2001). Histone H3 phosphorylation is coupled to poly-(ADP-ribosylation) during reactive oxygen species-induced cell death in renal proximal tubular epithelial cells. *Molecular Pharmacology* 60: 394-402.
- Tikoo K, Meena R, Kabra D, & Gaikwad A (2008). Change in post-translational modifications of histone H3, heat-shock protein-27 and MAP kinase p38 expression by curcumin in streptozotocin-induced type I diabetic nephropathy. *British Journal of Pharmacology* 153: 1225-1231.
- Tomita M, Asada M, Asada N, Nakamura J, Oguchi A, Higashi AY, *et al.* (2013). Bmp7 maintains undifferentiated kidney progenitor population and determines nephron numbers at birth. *PloS one* 8.
- Touyz RM, & Schiffrin EL (2000). Signal transduction mechanisms mediating the physiological and pathophysiological actions of angiotensin II in vascular smooth muscle cells. *Pharmacological Reviews* 52: 639-672.
- Tracy CM, Warren JS, Szulik M, Wang L, Garcia J, Makaju A, *et al.* (2018). The Smyd family of methyltransferases: Role in cardiac and skeletal muscle physiology and pathology. *Current Opinion in Physiology* 1: 140-152.
- Tracz M, Juncos J, Croatt A, Ackerman A, Grande JP, Knutson KL, *et al.* (2007). Deficiency of heme oxygenase-1 impairs renal hemodynamics and exaggerates systemic inflammatory responses to renal ischemia. *Kidney International* 72: 1073-1080.

- Tsai H-H, Yen R-F, Lin C-L, & Kao C-H (2017). Increased risk of dementia in patients hospitalized with acute kidney injury: A nationwide population-based cohort study. PloS one 12.
- TSAO N, Hsu H-P, WU C-M, Liu C-C, & LEI H-Y (2001). Tumour necrosis factor- α causes an increase in blood-brain barrier permeability during sepsis. Journal of Medical Microbiology 50: 812-821.
- Turner AJ, Tipnis SR, Guy JL, Rice GI, & Hooper NM (2002). ACEH/ACE2 is a novel mammalian metallocarboxypeptidase and a homologue of angiotensin-converting enzyme insensitive to ACE inhibitors. Canadian Journal of Physiology and Pharmacology 80: 346-353.
- Uchino S, Kellum JA, Bellomo R, Doig GS, Morimatsu H, Morgera S, *et al.* (2005). Acute renal failure in critically ill patients: a multinational, multicenter study. Jama 294: 813-818.
- Vaidya VS, Ferguson MA, & Bonventre JV (2008). Biomarkers of acute kidney injury. Annual Review of Pharmacology and Toxicology 48: 463-493.
- Van Beneden K, Geers C, Pauwels M, Mannaerts I, Verbeelen D, van Grunsven LA, *et al.* (2011). Valproic acid attenuates proteinuria and kidney injury. Journal of the American Society of Nephrology 22: 1863-1875.
- Varagic J, Ahmad S, Nagata S, & Ferrario CM (2014). ACE2: angiotensin II/angiotensin-(1-7) balance in cardiac and renal injury. Current Hypertension Reports 16: 1-9.
- Velkoska E, Patel SK, Griggs K, Pickering RJ, Tikellis C, & Burrell LM (2015). Short-Term Treatment with Diminazene Aceturate Ameliorates the Reduction in Kidney ACE2 Activity in Rats with Subtotal Nephrectomy. PloS One 10: e0118758.
- Venkatachalam MA, Weinberg JM, Kriz W, & Bidani AK (2015). Failed Tubule Recovery, AKI-CKD Transition, and Kidney Disease Progression. Journal of the American Society of Nephrology : JASN 26: 1765-1776.

- Verma L, Agrawal D, & Jain NS (2018). Enhanced central histaminergic transmission attenuates compulsive-like behavior in mice. *Neuropharmacology* 138: 106-117.
- Vickers C, Hales P, Kaushik V, Dick L, Gavin J, Tang J, *et al.* (2002). Hydrolysis of biological peptides by human angiotensin-converting enzyme-related carboxypeptidase. *The Journal of Biological Chemistry* 277: 14838-14843.
- Vilander LM, Kaunisto MA, & Pettilä V (2015). Genetic predisposition to acute kidney injury—a systematic review. *BMC Nephrology* 16: 197.
- Vilay AM, Churchwell MD, & Mueller BA (2008). Clinical review: drug metabolism and nonrenal clearance in acute kidney injury. *Critical Care* 12: 235.
- Villanueva S, Céspedes C, & Vio CP (2006). Ischemic acute renal failure induces the expression of a wide range of nephrogenic proteins. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* 290: R861-R870.
- Villar AJ, & Pedersen RA (1994). Parental imprinting of the Mas protooncogene in mouse. *Nature Genetics* 8: 373-379.
- Villela D, Leonhardt J, Patel N, Joseph J, Kirsch S, Hallberg A, *et al.* (2015). Angiotensin type 2 receptor (AT2R) and receptor Mas: a complex liaison. *Clinical Science* 128: 227-234.
- Villela DC, Passos-Silva DG, & Santos RAS (2014). Alamandine: a new member of the angiotensin family. *Current Opinion in Nephrology and Hypertension* 23: 130-134.
- Villeneuve LM, Reddy MA, Lanting LL, Wang M, Meng L, & Natarajan R (2008). Epigenetic histone H3 lysine 9 methylation in metabolic memory and inflammatory phenotype of vascular smooth muscle cells in diabetes. *Proceedings of the National Academy of Sciences* 105: 9047-9052.
- Waddington CH (1942). The epigenotype. *International Journal of Epidemiology* 41:10–131.

- Wagner T, & Jung M (2012). New lysine methyltransferase drug targets in cancer. *Nature Biotechnology* 30: 622-623.
- Wan Y, Wallinder C, Plouffe B, Beaudry H, Mahalingam A, Wu X, *et al.* (2004). Design, synthesis, and biological evaluation of the first selective nonpeptide AT2 receptor agonist. *Journal of Medicinal Chemistry* 47: 5995-6008.
- Wang J, Yan W, Peng X, Jiang Y, He L, Peng Y, *et al.* (2018). Functional role of SUV39H1 in human renal tubular epithelial cells under high-glucose ambiance. *Inflammation* 41: 1-10.
- Wang L, de Kloet AD, Pati D, Hiller H, Smith JA, Pioquinto DJ, *et al.* (2016). Increasing brain angiotensin converting enzyme 2 activity decreases anxiety-like behavior in male mice by activating central Mas receptors. *Neuropharmacology* 105: 114-123.
- Wang W, Koka V, & Lan HY (2005). Transforming growth factor- β and Smad signalling in kidney diseases. *Nephrology* 10: 48-56.
- Wei J, Yang F, Gong C, Shi X, & Wang G (2019). Protective effect of daidzein against streptozotocin-induced Alzheimer's disease via improving cognitive dysfunction and oxidative stress in rat model. *Journal of Biochemical and Molecular Toxicology*: e22319.
- Wei L, Clauser E, Alhenc-Gelas F, & Corvol P (1992). The two homologous domains of human angiotensin I-converting enzyme interact differently with competitive inhibitors. *Journal of Biological Chemistry* 267: 13398-13405.
- Williams P, Lopez H, Britt D, Chan C, Ezrin A, & Hottendorf R (1997). Characterization of renal ischemia-reperfusion injury in rats. *Journal of Pharmacological and Toxicological Methods* 37: 1-7.
- Wong M, Polly P, & Liu T (2015). The histone methyltransferase DOT1L: regulatory functions and a cancer therapy target. *American Journal of Cancer Research* 5: 2823-2837.
- Wong MKS (2016). Subchapter 29A - Renin. In *Handbook of Hormones*. eds Takei Y., Ando H., & Tsutsui K. Academic Press: San Diego, pp 255-e229A-252.

- Woo K-T, Choong H-L, Wong K-S, Tan H-K, Foo M, Stephanie F-C, *et al.* (2013). A retrospective Aliskiren and Losartan study in non-diabetic chronic kidney disease. *World Journal of Nephrology* 2: 129-133.
- Wu C-C, Wang I-F, Chiang P-M, Wang L-C, Shen C-KJ, & Tsai K-J (2017). G-CSF-mobilized bone marrow mesenchymal stem cells replenish neural lineages in Alzheimer's disease mice via CXCR4/SDF-1 chemotaxis. *Molecular Neurobiology* 54: 6198-6212.
- Wu V-C, Wu C-H, Huang T-M, Wang C-Y, Lai C-F, Shiao C-C, *et al.* (2014). Long-term risk of coronary events after AKI. *Journal of the American Society of Nephrology* 25: 595-605.
- Wu YS, Chen Y-T, Bao Y-T, Li Z-M, Zhou X-J, He J-N, *et al.* (2016). Identification and verification of potential therapeutic target genes in berberine-treated zucker diabetic fatty rats through bioinformatics analysis. *PloS one* 11: 0166378.
- Xia H, & Lazartigues E (2010). Angiotensin-converting enzyme 2: central regulator for cardiovascular function. *Current Hypertension Reports* 12: 170-175.
- Xiao F, Hiremath S, Knoll G, Zimpelmann J, Srivaratharajah K, Jadhav D, *et al.* (2012). Increased urinary angiotensin-converting enzyme 2 in renal transplant patients with diabetes. *PLoS One* 7: e37649.
- Xu P, Costa-Goncalves AC, Todiras M, Rabelo LA, Sampaio WO, Moura MM, *et al.* (2008). Endothelial dysfunction and elevated blood pressure in MAS gene-deleted mice. *Hypertension* 51: 574-580.
- Yacoub R, & Campbell KN (2015). Inhibition of RAS in diabetic nephropathy. *Int J Nephrol Renovasc Dis* 8: 29-40.
- Yan W, Herman JG, & Guo M (2016). Epigenome-based personalized medicine in human cancer. *Epigenomics* 8: 119-133.

- Yang G, Weng X, Zhao Y, Zhang X, Hu Y, Dai X, *et al.* (2017a). The histone H3K9 methyltransferase SUV39H links SIRT1 repression to myocardial infarction. *Nature Communications* 8: 1-15.
- Yang G, Zhang X, Weng X, Liang P, Dai X, Zeng S, *et al.* (2017b). SUV39H1 mediated SIRT1 trans-repression contributes to cardiac ischemia–reperfusion injury. *Basic Research in Cardiology* 112: 1-14.
- Yang L (2016). Acute kidney injury in Asia. *Kidney Diseases* 2: 95-102.
- Yang Xh, Wang Yh, Wang Jj, Liu Yc, Deng W, Qin C, *et al.* (2012). Role of angiotensin-converting enzyme (ACE and ACE2) imbalance on tourniquet-induced remote kidney injury in a mouse hindlimb ischemia–reperfusion model. *Peptides* 36: 60-70.
- Yang Y, Yin B, Lv L, Wang Z, He J, Chen Z, *et al.* (2017c). Gastroprotective effect of aucubin against ethanol-induced gastric mucosal injury in mice. *Life Sciences* 189: 44-51.
- Yap SC, & Lee HT (2012). Acute Kidney Injury and Extrarenal Organ DysfunctionNew Concepts and Experimental Evidence. *Anesthesiology: The Journal of the American Society of Anesthesiologists* 116: 1139-1148.
- Yu C, & Zhuang S (2019). Histone Methyltransferases as Therapeutic Targets for Kidney Diseases. *Frontiers in Pharmacology* 10: 1-9.
- Yu H, Lin Q, Wang Y, He Y, Fu S, Jiang H, *et al.* (2013). Inhibition of H3K9 methyltransferases G9a/GLP prevents ototoxicity and ongoing hair cell death. *Cell Death & Disease* 4: e506-e506.
- Yu SM-W, & Bonventre JV (2018). Acute kidney injury and progression of diabetic kidney disease. *Advances in Chronic Kidney Disease* 25: 166-180.
- Yuan H, & Marmorstein R (2013). Histone acetyltransferases: Rising ancient counterparts to protein kinases. *Biopolymers* 99: 98-111.

- Yvan-Charvet L, Even P, Bloch-Faure M, Guerre-Millo M, Moustaid-Moussa N, Ferre P, *et al.* (2005). Deletion of the angiotensin type 2 receptor (AT2R) reduces adipose cell size and protects from diet-induced obesity and insulin resistance. *Diabetes* 54: 991-999.
- Zacharias N, Sailhamer EA, Li Y, Liu B, Butt MU, Shuja F, *et al.* (2011). Histone deacetylase inhibitors prevent apoptosis following lethal hemorrhagic shock in rodent kidney cells. *Resuscitation* 82: 105-109.
- Zager R, Johnson A, Hanson S, & Lund S (2006). Acute nephrotoxic and obstructive injury primes the kidney to endotoxin-driven cytokine/chemokine production. *Kidney International* 69: 1181-1188.
- Zager R, Johnson A, & Lund S (2007). ‘Endotoxin tolerance’: TNF- α hyper-reactivity and tubular cytoresistance in a renal cholesterol loading state. *Kidney International* 71: 496-503.
- Zager R, Johnson AC, Hanson SY, & Lund S (2005). Ischemic proximal tubular injury primes mice to endotoxin-induced TNF- α generation and systemic release. *American Journal of Physiology-Renal Physiology* 289: F289-F297.
- Zager RA, Johnson AC, & Becker K (2011). Acute unilateral ischemic renal injury induces progressive renal inflammation, lipid accumulation, histone modification, and “end-stage” kidney disease. *American Journal of Physiology-Renal Physiology* 301: F1334-F1345.
- Zager RA, Johnson AC, & Frostad KB (2014). Acute hepatic ischemic-reperfusion injury induces a renal cortical “stress response,” renal “cytoresistance,” and an endotoxin hyperresponsive state. *American Journal of Physiology-Renal Physiology* 307: F856-F868.
- Zhang G, Wang Q, Zhou Q, Wang R, Xu M, Wang H, *et al.* (2016). Protective effect of tempol on acute kidney injury through PI3K/Akt/Nrf2 signaling pathway. *Kidney and Blood Pressure Research* 41: 129-138.

- Zhang H, & Sun S-C (2015). NF-κB in inflammation and renal diseases. *Cell & Bioscience* 5: 1-12.
- Zhang J, He P, Xi Y, Geng M, Chen Y, & Ding J (2015). Down-regulation of G9a triggers DNA damage response and inhibits colorectal cancer cells proliferation. *Oncotarget* 6: 2917-2927.
- Zhang Q, Raoof M, Chen Y, Sumi Y, Sursal T, Junger W, *et al.* (2010). Circulating mitochondrial DAMPs cause inflammatory responses to injury. *Nature* 464: 104-107.
- Zhang Y (2003). Transcriptional regulation by histone ubiquitination and deubiquitination. *Genes & Development* 17: 2733-2740.
- Zhang Y, Griffin K, Mondal N, & Parvin JD (2004). Phosphorylation of histone H2A inhibits transcription on chromatin templates. *Journal of Biological Chemistry* 279: 21866-21872.
- Zhao H, Huang H, Alam A, Chen Q, Suen KC, Cui J, *et al.* (2018). VEGF mitigates histone-induced pyroptosis in the remote liver injury associated with renal allograft ischemia–reperfusion injury in rats. *American Journal of Transplantation* 18: 1890-1903.
- Zhou Q, Liu K, Wu H, Chen L, Pouranan V, Yuan M, *et al.* (2012). Spironolactone rescues Dot1a-Af9-mediated repression of endothelin-1 and improves kidney injury in streptozotocin-induced diabetic rats. *PLoS One* 7: e47360.
- Zhou X, Ono H, Ono Y, & Frohlich ED (2004). Aldosterone antagonism ameliorates proteinuria and nephrosclerosis independent of glomerular dynamics in L-NAME/SHR model. *American Journal of Nephrology* 24: 242-249.
- Zhou X, Xiong C, Tolbert E, Zhao TC, Bayliss G, & Zhuang S (2018). Targeting histone methyltransferase enhancer of zeste homolog-2 inhibits renal epithelial–mesenchymal transition and attenuates renal fibrosis. *The FASEB Journal* 32: 5976-5989.
- Zhou X, Zang X, Ponnusamy M, Masucci MV, Tolbert E, Gong R, *et al.* (2016). Enhancer of zeste homolog 2 inhibition attenuates renal fibrosis by maintaining Smad7 and

- phosphatase and tensin homolog expression. *Journal of the American Society of Nephrology* 27: 2092-2108.
- Zhu B, Zheng Y, Pham A-D, Mandal SS, Erdjument-Bromage H, Tempst P, *et al.* (2005). Monoubiquitination of human histone H2B: the factors involved and their roles in HOX gene regulation. *Molecular Cell* 20: 601-611.
- Zhuang S (2018). Epigenetic targeting for acute kidney injury. *Nephrology* 23: 21-25.
- Zilberman-Itskovich S, Abu-Hamad R, Zarura R, Sova M, Hachmo Y, Stark M, *et al.* (2019). Human mesenchymal stromal cells ameliorate complement induced inflammatory cascade and improve renal functions in a rat model of ischemia-reperfusion induced acute kidney injury. *PLoS One* 14: e0222354.
- Ziypak T, Halici Z, Alkan E, Akpinar E, Polat B, Adanur S, *et al.* (2015). Renoprotective effect of aliskiren on renal ischemia/reperfusion injury in rats: electron microscopy and molecular study. *Renal Failure* 37: 343-354.
- Zou XJ, Yang L, & Yao SL (2012). Endoplasmic reticulum stress and C/EBP homologous protein-induced Bax translocation are involved in angiotensin II-induced apoptosis in cultured neonatal rat cardiomyocytes. *Experimental Biology and Medicine* 237: 1341-1349.
- Zuk A, & Bonventre JV (2016). Acute kidney injury. *Annual Review of Medicine* 67: 293-307.