

List of Figures

1.1	Numerical solution of (1.2) using uniform mesh with $\varepsilon = 10^{-4}$ and $N = 256$	12
1.2	Numerical solution of (1.2) using uniform mesh with $\varepsilon = 10^{-4}$ and $N = 2^{13}$	13
2.1	Numerical solution profiles for Example 2.6.1 for different values of ε (a) $\varepsilon = 1$ (b) $\varepsilon = 2^{-4}$ (c) $\varepsilon = 2^{-8}$ and (d) $\varepsilon = 2^{-12}$	48
2.2	The log-log error plot for Example 2.6.1.	49
2.3	Numerical solution for Example 2.6.1 for different values of ε and t (a) $\varepsilon = 1$ (b) $\varepsilon = 0.1$ (c) $\varepsilon = 0.01$ and (d) $\varepsilon = 0.005$	50
2.4	Numerical solution profiles for Example 2.6.2 for different values of ε (a) $\varepsilon = 1$ (b) $\varepsilon = 2^{-4}$ (c) $\varepsilon = 2^{-8}$ and (d) $\varepsilon = 2^{-12}$	52
2.5	The log-log error plot for Example 2.6.2.	53
2.6	Numerical solution for Example 2.6.2 for different values of ε and t (a) $\varepsilon = 1$ (b) $\varepsilon = 0.1$ (c) $\varepsilon = 0.01$ and (d) $\varepsilon = 0.005$	54
3.1	Surface plots for Example 3.6.1 for (a) $\varepsilon = 1$ (b) $\varepsilon = 2^{-4}$ (c) $\varepsilon = 2^{-8}$ and (d) $\varepsilon = 2^{-12}$	78
3.2	Numerical solution for Example 3.6.1 at different time levels for (a) $\varepsilon = 1$ (b) $\varepsilon = 0.1$ (c) $\varepsilon = 0.01$ and (d) $\varepsilon = 0.001$	79
3.3	Absolute errors profile for Example 3.6.1 for $\varepsilon = 0.01$	80
3.4	Surface plots for Example 3.6.2 for (a) $\varepsilon = 1$ (b) $\varepsilon = 2^{-4}$ (c) $\varepsilon = 2^{-8}$ (d) $\varepsilon = 2^{-12}$	82
3.5	Numerical solution for Example 3.6.2 at different time levels for (a) $\varepsilon = 1$ (b) $\varepsilon = 0.1$ (c) $\varepsilon = 0.01$ and (d) $\varepsilon = 0.001$	83
3.6	Absolute errors profile for Example 3.6.2 for $\varepsilon = 0.01$	84
4.1	Numerical solution profiles for Example 4.5.1 (a) $\varepsilon = 1$ (b) $\varepsilon = 2^{-4}$ (c) $\varepsilon = 2^{-8}$ and (d) $\varepsilon = 2^{-12}$	102

4.2	Numerical solution profiles for Example 4.5.1 for different time levels (a) $\varepsilon = 1$ (b) $\varepsilon = 0.1$ (c) $\varepsilon = 0.01$ and (d) $\varepsilon = 0.001$	103
4.3	Numerical solution profiles for Example 4.5.2 (a) $\varepsilon = 1$ (b) $\varepsilon = 2^{-4}$ (c) $\varepsilon = 2^{-8}$ and (d) $\varepsilon = 2^{-12}$	105
4.4	Numerical solution profiles for Example 4.5.2 for different time levels (a) $\varepsilon = 1$ (b) $\varepsilon = 0.1$ (c) $\varepsilon = 0.01$ and (d) $\varepsilon = 0.001$	106
5.1	Numerical solution profiles for Example 5.5.1 for (a) $\varepsilon = 1, p = 1$ (b) $\varepsilon = 2^{-6}, p = 3$ (c) $\varepsilon = 2^{-12}, p = 5$ and (d) $\varepsilon = 2^{-18}, p = 7$	124
5.2	Numerical solution profiles for $p = 2$ at different time levels for Ex- ample 5.5.1 for (a) $\varepsilon = 1$ (b) $\varepsilon = 0.1$ (c) $\varepsilon = 0.01$ and (d) $\varepsilon = 0.001$. . .	125
5.3	Numerical solution profiles for Example 5.5.2 for (a) $\varepsilon = 1, p = 1$ (b) $\varepsilon = 2^{-6}, p = 3$ (c) $\varepsilon = 2^{-12}, p = 5$ and (d) $\varepsilon = 2^{-18}, p = 7$	128
5.4	Numerical solution profiles for $p = 2$ at different time levels for Ex- ample 5.5.2 for (a) $\varepsilon = 1$ (b) $\varepsilon = 0.1$ (c) $\varepsilon = 0.01$ and (d) $\varepsilon = 0.001$. . .	129
6.1	Numerical solution profiles for Example 6.5.1 for (a) $\varepsilon_1 = 2^{-15}, \varepsilon_2 =$ 2^{-6} (b) $\varepsilon_1 = 2^{-10}, \varepsilon_2 = 2^{-8}$ (c) $\varepsilon_1 = 2^{-17}, \varepsilon_2 = 2^{-7}$ and (d) $\varepsilon_1 =$ $2^{-14}, \varepsilon_2 = 2^{-18}$	152
6.2	Numerical solution profiles for Example 6.5.1 for $\varepsilon_1 = 2^{-14}, \varepsilon_2 = 2^{-18}$	153
6.3	Numerical solution profiles for Example 6.5.2 (viewed from right) for (a) $\varepsilon_1 = 2^{-15}, \varepsilon_2 = 2^{-6}$ (b) $\varepsilon_1 = 2^{-10}, \varepsilon_2 = 2^{-8}$ (c) $\varepsilon_1 = 2^{-17}, \varepsilon_2 =$ 2^{-7} and (d) $\varepsilon_1 = 2^{-14}, \varepsilon_2 = 2^{-18}$	156
6.4	Numerical solution profiles for Example 6.5.2 (viewed from left) for (a) $\varepsilon_1 = 2^{-15}, \varepsilon_2 = 2^{-6}$ (b) $\varepsilon_1 = 2^{-10}, \varepsilon_2 = 2^{-8}$ (c) $\varepsilon_1 = 2^{-17}, \varepsilon_2 =$ 2^{-7} and (d) $\varepsilon_1 = 2^{-14}, \varepsilon_2 = 2^{-18}$	157
6.5	Numerical solution profiles for Example 6.5.2 for $\varepsilon_1 = 2^{-14}, \varepsilon_2 = 2^{-18}$	158
7.1	Numerical solution profiles for Example 7.5.1.	185
7.2	Numerical solution profiles for Example 7.5.1.	185
7.3	Numerical solution profiles for $\varepsilon_1 = 2^{-10}, \varepsilon_2 = 2^{-4}$ by taking contin- uous source terms.	186
7.4	Numerical solution profiles for Example 7.5.2.	188
7.5	Numerical solution profiles for Example 7.5.2.	188

List of Tables

2.1	$E_\varepsilon^{N,M}$, $E^{N,M}$, $p_\varepsilon^{N,M}$ and $p^{N,M}$ for Example 2.6.1.	47
2.2	$E_\varepsilon^{N,M}$, $E^{N,M}$, $p_\varepsilon^{N,M}$ and $p^{N,M}$ for Example 2.6.2.	51
3.1	$e_{N,\Delta t}^\varepsilon$, $e_{N,\Delta t}$, $\rho_{N,\Delta t}^\varepsilon$ and $\rho_{N,\Delta t}$ for Example 3.6.1.	77
3.2	$e_{N,\Delta t}^\varepsilon$, $e_{N,\Delta t}$, $\rho_{N,\Delta t}^\varepsilon$ and $\rho_{N,\Delta t}$ for Example 3.6.2.	81
4.1	$e_\varepsilon^{N,M}$, $e^{N,M}$, $p_\varepsilon^{N,M}$ and $p^{N,M}$ for Example 4.5.1.	101
4.2	$e_\varepsilon^{N,M}$, $e^{N,M}$, $p_\varepsilon^{N,M}$ and $p^{N,M}$ for Example 4.5.2.	104
5.1	$e_\varepsilon^{N,M}$, $e^{N,M}$, $q_\varepsilon^{N,M}$ and $q^{N,M}$ for Example 5.5.1 for $p = 1$	122
5.2	$e_\varepsilon^{N,M}$ and $q_\varepsilon^{N,M}$ for Example 5.5.1 for $\varepsilon = 2^{-10}$ and different values of p	123
5.3	$e_\varepsilon^{N,M}$, $e^{N,M}$, $q_\varepsilon^{N,M}$ and $q^{N,M}$ for Example 5.5.2 for $p = 1$	126
5.4	$e_\varepsilon^{N,M}$ and $q_\varepsilon^{N,M}$ for Example 5.5.2 for $\varepsilon = 2^{-10}$ and different values of p	127
6.1	Computational results for Example 6.5.1 for $\varepsilon_1 = 2^{-10}$ and different values of ε_2	150
6.2	Computational results for Example 6.5.1 for $\varepsilon_2 = 2^{-12}$ and different values of ε_1	151
6.3	Computational results for Example 6.5.2 for $\varepsilon_1 = 2^{-10}$ and different values of ε_2	154
6.4	Computational results for Example 6.5.2 for $\varepsilon_2 = 2^{-12}$ and different values of ε_1	155
7.1	$e_{\varepsilon_1,\varepsilon_2}^{N,M}$ and $\rho_{\varepsilon_1,\varepsilon_2}^{N,M}$ for Example 7.5.1 for $\varepsilon_1 = 2^{-6}$	181
7.2	$e_{\varepsilon_1,\varepsilon_2}^{N,M}$ and $\rho_{\varepsilon_1,\varepsilon_2}^{N,M}$ for Example 7.5.1 for $\varepsilon_2 = 2^{-8}$	182
7.3	$e_{\varepsilon_1,\varepsilon_2}^{N,M}$ and $\rho_{\varepsilon_1,\varepsilon_2}^{N,M}$ for Example 7.5.1 by taking $N = 256$	183
7.4	Comparison of $e_{\varepsilon_1,\varepsilon_2}^{N,M}$ for Example 7.5.1 for $\varepsilon_2 = 2^{-4}$	184
7.5	$e_{\varepsilon_1,\varepsilon_2}^{N,M}$ and $\rho_{\varepsilon_1,\varepsilon_2}^{N,M}$ for Example 7.5.2 for $\varepsilon_1 = 2^{-15}$	186
7.6	$e_{\varepsilon_1,\varepsilon_2}^{N,M}$ and $\rho_{\varepsilon_1,\varepsilon_2}^{N,M}$ for Example 7.5.2 for $\varepsilon_2 = 2^{-16}$	187